

# REDOX REACTION & EQUIVALENT CONCEPT

## OXIDATION & REDUCTION :

Let us do a comparative study of oxidation and Reduction ;

Oxidation		Reduction	
(1)	Addition of oxygen <b>e.g.</b> $2\text{Mg} + \text{O}_2 \rightarrow 2\text{MgO}$	(i)	Removal of oxygen <b>e.g.</b> $\text{CuO} + \text{C} \rightarrow \text{Cu} + \text{CO}$
(2)	Removal of Hydrogen <b>e.g.</b> $\text{H}_2\text{S} + \text{Cl}_2 \rightarrow 2\text{HCl} + \text{S}$	(ii)	Addition of Hydrogen <b>e.g.</b> $\text{S} + \text{H}_2 \rightarrow \text{H}_2\text{S}$
(3)	Increase in positive charge <b>e.g.</b> $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + e^-$	(iii)	Decrease in positive charge <b>e.g.</b> $\text{Fe}^{3+} + e^- \rightarrow \text{Fe}^{2+}$
(4)	Increase in oxidation number (+2)      (+4) $\text{SnCl}_2 \rightarrow \text{SnCl}_4$	(iv)	Decrease in oxidation number (+7)      (+2) $\text{MnO}_4^- \rightarrow \text{Mn}^{2+}$
(5)	Removal of electron <b>e.g.</b> $\text{Sn}^{2+} \rightarrow \text{Sn}^{4+} + 2e^-$	(v)	Addition of electron <b>e.g.</b> $\text{Fe}^{3+} + e^- \rightarrow \text{Fe}^{2+}$

## Oxidation Number :

- It is an imaginary or apparent charge gained by an element when it goes from its elemental free state to combined state in molecules.
- It is calculated on basis of a arbitrary set of rules.
- It is a relative charge in a particular bonded state.

## Rules governing oxidation number :

The following rules are helpful in calculating oxidation number of the elements in their different compounds. It is remember that the basis of these rule is the electronegativity of the element.

### Fluorine atom :

Fluorine is most electronegativity atom (known). It always has oxidation no. equal to -1 in all its compounds.

### Oxygen atom :

In general and as well as in its oxides, oxygen atom has oxidation number equal to -2.

- In case of :
- peroxide (e.g.  $\text{H}_2\text{O}_2$ ,  $\text{Na}_2\text{O}_2$ ) is -1
  - super oxide (e.g.  $\text{KO}_2$ ) is  $-\frac{1}{2}$
  - ozonide ( $\text{KO}_3$ ) is  $-\frac{1}{3}$
  - oxygen fluoride  $\text{OF}_2$  is +2 & in  $\text{O}_2\text{F}_2$  is +1


### Hydrogen atom :

In general, H atom has oxidation number equal to +1. But in metallic hydrides (e.g. NaH, KH) it is -1.

### Halogen atom :

In general, all halogen atom (Cl, Br, I) has oxidation number equal to -1.

But if halogen atom is attached with an more electronegative atom than halogen atom then it will show positive oxidation numbers.



◆ **Metals :**

(a) Alkali metal (Li, Na, K, Rb, ..... ) always have oxidation number +1.

(b) Alkaline earth metal (Be, Mg, Ca ..... ) always have oxidation number +2.

**Note :** Metal may have negative or zero oxidation number.

(c) Aluminium always have +3 oxidation number

◆ Oxidation number of an element in free state or in allotropic forms is always zero.

e.g.  $\overset{0}{\text{O}}_2, \overset{0}{\text{S}}_8, \overset{0}{\text{P}}_4, \overset{0}{\text{O}}_3$

◆ Sum of the charges of elements in a molecule is zero.

◆ Sum of the charges of all elements in an ions is equal to the charge on the ion.

◆ If the group no. of an element in periodic table is n then its oxidation number may vary from n to n – 8 (but it is mainly applicable in p-block elements)

e.g. N-atom belongs to v group in the periodic table therefore as per rule its oxidation number may vary from  
 $\overset{-3}{\text{N}}\text{H}_3, \overset{+2}{\text{N}}\text{O}, \overset{+3}{\text{N}}_2\text{O}_3, \overset{+4}{\text{N}}\text{O}_2, \overset{+5}{\text{N}}_2\text{O}_5$   
 -3 to +5 (NH<sub>3</sub>, NO, N<sub>2</sub>O<sub>3</sub>, NO<sub>2</sub>, N<sub>2</sub>O<sub>5</sub>)

**Calculation of average oxidation number :**

**Solved Examples :**

**Ex.** Calculate oxidation number of underlined element Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> :

**Sol.** Let oxidation number of S-atom is x. Now work accordingly with the rules given before.

$$(+1) \quad 2 + (x) \quad 2 + (-2) \quad 3 = 0$$

$$x = + 2$$

**Ex.** Na<sub>2</sub>S<sub>4</sub>O<sub>6</sub> :

**Sol.** Let oxidation number of S-atom is x

$$(+1) \quad 2 + (x) \quad 2 + (-2) \quad 6 = 0$$

$$x = + 2.5$$

◆ It's important to note here that Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> have two S-atom and there are four S-atom in Na<sub>2</sub>S<sub>4</sub>O<sub>6</sub> but none sulphur atom in both the compound have +2 or +2.5 oxidation number, it is the average charge (O. No.) Which reside on each sulphur atom therefore we should work to calculate the individual oxidation number of each sulphur atom in these compound.

**Ex.** Calculate the O.S. of all the atoms in the following species :

(i) ClO<sup>-</sup>, (ii) NO<sub>2</sub><sup>-</sup>, (iii) NO<sub>3</sub><sup>-</sup> (iv) CCl<sub>4</sub> (v) K<sub>2</sub>CrO<sub>4</sub> and (vi) KMnO<sub>4</sub>

**Sol.** (i) In ClO<sup>-</sup>, the net charge on the species is -1 and therefore the sum of the oxidation states of Cl and O must be equal to -1. Oxygen will have an O.S. of -2 and if the O.S. of Cl is assumed to be 'x' then x - 2 should be equal to -1.

$$x \text{ is } + 1$$

(ii) NO<sub>2</sub><sup>-</sup> : (2 -2) + x = -1 (where 'x' is O.S. of N)

$$\therefore x = +3$$

(iii) NO<sub>3</sub><sup>-</sup> : x + (3 -2) = -1 (where 'x' is O.S. of N)

$$x = +5$$

(iv) In CCl<sub>4</sub>, Cl has an O.S. of -1

$$x + 4 \quad -1 = 0$$


$$\therefore x = +4$$

(where 'x' is O.S. of C)

- (v)  $\text{K}_2\text{CrO}_4$  : K has O.S. of +1 and O has O.S. of -2 and let Cr has O.S. 'x' then,  
 $2(+1) + x + 4(-2) = 0$   
 $\therefore x = +6$
- (vi)  $\text{KMnO}_4$  :  $+1 + x + (4 \times -2) = 0$   
 $\therefore x = +7$  (where x is O.S. of Mn).

#### ❑ MISCELLANEOUS EXAMPLES :

In order to determine the exact or individual oxidation number we need to take help from the structures of the molecules. Some special cases are discussed as follows :




From the structure it is evident that in  $\text{CrO}_5$  there are two peroxide linkages and one double bond.

The contribution of each peroxide linkage is -2. Let the O.N. of Cr is x.

$$\therefore x + (-2)2 + (-2) = 0 \text{ or } x = 6$$

$$\therefore \text{O.N. of Cr} = +6 \text{ Ans.}$$



From the structure, it is evident that in  $\text{H}_2\text{SO}_5$ , there are one peroxide linkage, two sulphur - oxygen double bond and one OH group. Let the O.N. of S = x.

$$\therefore +1 - 2 + x + (-2)2 + (-2) + 1 = 0$$

$$\text{or } x + 2 - 8 = 0$$

$$\text{or } x - 6 = 0$$


$$\text{or } x = 6$$

$$\therefore \text{O.N. of S in } \text{H}_2\text{SO}_5 \text{ is } +6 \text{ Ans}$$

#### ❑ Paradox of fractional oxidation number :


Fractional oxidation state is the average oxidation state of the element under examination and the structural parameters reveal that the element for whom fractional oxidation state is realised is actually present in different oxidation states. Structure of the species  $\text{C}_3\text{O}_2$ ,  $\text{Br}_3\text{O}_8$  and  $\text{S}_4\text{O}_6^{2-}$  reveal the following bonding situations.

- ◆ The element marked with asterisk in each species is exhibiting the different oxidation state (oxidation number) from rest of the atoms of the same element in each of the species. This reveals that in  $\text{C}_3\text{O}_2$ , two carbon atoms are present in +2 oxidation state each whereas the third one is present in zero oxidation state and the average is  $4/3$ . However the realistic picture is +2 for two terminal carbons and zero for the middle carbon.




Structure of  $\text{C}_3\text{O}_2$   
 (Carbon suboxide)

- ◆ Likewise in  $\text{Br}_3\text{O}_8$ , each of the two terminal bromine atoms are present in +6 oxidation state and the middle bromine is present in +4 oxidation state. Once again the average, that is different from reality is  $16/3$ .



Structure of  $\text{Br}_3\text{O}_8$  (tribromooctaoxide)

- ◆ In the same fashion, in the species  $\text{S}_4\text{O}_6^{2-}$ , is 2.5, whereas the reality being +5, 0, 0 and +5 oxidation number respectively for each sulphur.



Structure of  $\text{S}_4\text{O}_6^{2-}$  (tetrathionate ion)

## ❑ OXIDISING AND REDUCING AGENT :

### ◆ Oxidising agent or Oxidant :

Oxidising agents are those compound which can oxidise others and reduced itself during the chemical reaction. Those reagents whose O.N. decrease or which gain electrons in a redox reaction are termed as oxidants

e.g.  $\text{KMnO}_4$ ,  $\text{K}_2\text{Cr}_2\text{O}_7$ ,  $\text{HNO}_3$ , conc.  $\text{H}_2\text{SO}_4$  etc, are powerful oxidising agents.

### ◆ Reducing agent or Reductant :


Reducing agents are those compound which can reduce others and oxidise itself during the chemical reaction. Those reagents whose O.N. increase or which loses electrons in a redox reaction are termed as reductants.

e.g.  $\text{KI}$ ,  $\text{Na}_2\text{S}_2\text{O}_3$  are powerful reducing agents.

**Note :** There are some compounds also which can work both oxidising agent and reducing agent.

e.g.  $\text{H}_2\text{O}_2$ ,  $\text{NO}_2^-$


## ❑ HOW TO IDENTIFY WHETHER A PARTICULAR SUBSTANCE IS AN OXIDISING OR REDUCING AGENT :



## ❑ REDOX REACTION :

A reaction in which oxidation and reduction simultaneously take place.




In all redox reactions the total increase in oxidation number must equal the total decrease in oxidation number.




### ◆ Disproportionations reactions :

A redox reaction in which a same element present in a particular compound in a definite oxidation state is oxidized as well as reduced simultaneously is a disproportionation reactions.


Disproportionation reactions are a special type of redox reactions. One of the reactants in a disproportionation reaction always contains **an element that can exist in at least three oxidation states**. The element in the form of reacting substance is in the intermediate oxidation state and both higher and lower oxidation states of that element are formed in the reaction.




◆ Consider following reactions :



$\text{KClO}_3$  plays a role of oxidant and reductant both. Because same element is not oxidised and reduced. Here, Cl present in  $\text{KClO}_3$  is reduced and O present in  $\text{KClO}_3$  is oxidized. So **its not a disproportion reaction** although it looks like one.






Nitrogen in this compound has  $-3$  and  $+3$  oxidation number so it is not a definite value, so its not a disproportion reaction. Its a example of **comproportionation reaction** which is a class of redox reaction in which a element from two different oxidation state gets converted into a single oxidation state.





Its a case of disproportionation reaction in which Cl is the atom disproportionating.


◆ List of some important disproportionation reaction :







**$\text{F}_2$  does not (can not) undergo disproportionation as it is the most electronegative element.**




◆ Reverse of disproportionation is called Comproportionation. In some of the disproportionation reactions by changing the medium (from acidic to basic or reverse) the reaction goes in backward direction and can be taken as an example of Comproportionation.



□ Some examples of redox reactions are :







⇒ If one of the half reactions does not take place, other half will also not take place. We can say oxidation and reduction go side by side.



In the above reaction, we can see that  $\text{Cl}_2$  has been oxidized as well as reduced. Such type of redox reaction is called **disproportionation** reaction.

□ **Some More examples :**



□ **BALANCING OF REDOX REACTION :**

All balanced equations must satisfy two criteria

1. **Atom balance (mass balance) :**

That is there should be the same number of atoms of each kind in reactant and products side.

2. **Charge balance :**

That is the sum of actual charges on both side of the equation must be equal

There are two methods for balancing the redox equations

(a) Oxidation - number change method

(b) Ion electron method or half cell method


(a) **Oxidation number change method :**

This method was given by Jonson. In a balanced redox reaction, total increase in oxidation number must be equal to total decreases in oxidation number. This equivalence provides the basis for balancing redox reactions.


The general procedure involves the following steps :

- (i) Select the atom in oxidising agent whose oxidation number decreases and indicate the gain of electrons.
- (ii) Select the atom in reducing agent whose oxidation number increases and write the loss of electrons.
- (iii) Now cross multiply i.e. multiply oxidising agent by the number of loss of electrons and reducing agent by number of gain of electrons.
- (iv) Balance the number of atoms on both sides whose oxidation numbers change in the reaction.
- (v) In order to balance oxygen atoms, add  $\text{H}_2\text{O}$  molecules to the side deficient in oxygen. Then balance the number of H atoms by adding  $\text{H}^+$  ions in the hydrogen.

**Ex.** Balance the following reaction by the oxidation number method :




**Sol.** Write the oxidation number of all the atoms.




There is change in oxidation number of Cu and N.



To make increase and decrease equal, eq. (2) is multiplied by 2.



Balancing nitrates ions, hydrogen and oxygen, the following equation is obtained.




This is the balanced equation.


**Ex.** Write the skeleton equation for each of the following processes and balance them by ion electron method :

- (i) Permagnet ion oxidizes oxalate ions in acidic medium to carbon dioxide and gets reduced itself to  $\text{Mn}^{2+}$  ions.
- (ii) Bromine and hydrogen peroxide react to give bromate ions and water.
- (iii) Chlorine reacts with base to form chlorate ion, chloride ion and water in acidic medium.

**Sol.** (i) The skeleton equation for the process :





◆ **Step (1)** : Indicating oxidation number :





◆ **Step (2)** : Writing oxidation and reduction half reaction :





◆ **Step (3)** : Adding electrons to make the difference in O.N.





◆ **Step (4)** : Balancing 'O' atom by adding  $\text{H}_2\text{O}$  molecules




◆ **Step (5)** : Balancing H atom by adding  $\text{H}^+$  ions




◆ **Step (6)** : Multiply the oxidation half reaction by 2 and reduction half reaction by 5 to equalize the electrons lost and gained and add the two half reactions.



(ii) The skeleton equation for the given process is



◆ **Step (1)** : Indicate the oxidation number of each atom





Thus, Br and O changes their oxidation numbers.



◆ **Step (2)** : Write the oxidation and reduction half reaction.





◆ **Step (3)** : Addition of electrons to make up for the difference in O.N.




◆ **Step (4)** : Balance 'O' atoms by adding  $\text{H}_2\text{O}$  molecules




◆ **Step (5)** : Equalize the electrons by multiplying the reduction half with 5 and add the two half reactions



(iii) The skeleton equation for the given process :




◆ **Step (1)** : Indicate the oxidation number of each atom





Thus, chlorine is the only element which undergoes the change in oxidation number. It decreases its oxidation number from 0 to -1 and also increases its oxidation number from 0 to +5.

◆ **Step (2)** : Write the oxidation and reduction half reactions





◆ **Step (3)** : Add electrons to make up for the difference in O.N.












- ◆ **Step (4) :** Balance O atoms by adding H<sub>2</sub>O molecules




- ◆ **Step (5) :** Since medium is basic, balance H atoms by adding H<sub>2</sub>O molecules to the side falling short of H atoms and equal number of OH<sup>-</sup> ions to the other side.



- ◆ **Step (6) :** Multiply the reduction half reaction by 5 and add two half reactions.




**Ex :** Balance the following reaction by the oxidation number method :



**Sol :** Write the oxidation number of all the atoms.


+7 -2



change in oxidation number has occurred in Mn and Fe.

+7

+2



.....(1)


(Decrement in oxidation no. by 5)



.....(2)

(Increment in oxidation no. by 1)

To make increase and decrease equal, eq. (2) is multiplied by 5.



To balance oxygen, 4H<sub>2</sub>O are added to R.H.S. and to balance hydrogen, 8H<sup>+</sup> are added to L.H.S.






This is the balanced equation.

**Ex.** Balance the following chemical reaction by oxidation number method and write their skeleton equation :


- Chloride ions reduce manganese dioxide to manganese (II) ions in acidic medium and get itself oxidized to chlorine gas.
- The nitrate ions in acidic medium oxidize magnesium to Mg<sup>2+</sup> ions but itself gets reduced to nitrous oxide.

**Sol.** (i) The skeleton equation for the given process is




- ◆ **Step (3) :** Equalize the increase/decrease in O.N. by multiplying


$\text{MnO}_2$  by 1 and  $\text{Cl}^-$  by 2



- ◆ **Step (4) :** Balance other atoms except H and O. Here they are all balanced.
- ◆ **Step (5) :** Balance O atoms by adding  $\text{H}_2\text{O}$  molecules to the side falling short of O atoms.



- ◆ **Step (6) :** Balance H atoms by adding  $\text{H}^+$  ions to the side falling short of H atoms.



- (ii) The skeleton equation for the given process is




- ◆ **Step (1) :**  $\overset{0}{\text{Mg}} + (\overset{+5}{\text{N}}\overset{-2}{\text{O}_3})^- \longrightarrow \overset{+2}{\text{Mg}} + \overset{+1}{\text{N}}\overset{-2}{\text{O}} + \overset{+1}{\text{H}}\overset{-2}{\text{O}}$

Multiply  $\text{NO}_3^-$  by 2 to equalize N atoms


- ◆ **Step (2) :**  $\overset{0}{\text{Mg}} + 2\overset{+5}{\text{N}}\overset{-2}{\text{O}_3} \longrightarrow \overset{+2}{\text{Mg}} + \overset{+1}{\text{N}}\overset{-2}{\text{O}} + \text{H}_2\text{O}$   

$\begin{array}{c} \text{O.N. increases by 2 per Mg atom} \\ \text{O.N. decreases by 4 per N atom} \end{array}$


- ◆ **Step (3) :** Equalize increase/decrease in O.N. by multiplying Mg by 4 and  $2\text{NO}_3^-$  by 1.



- ◆ **Step (4) :** Balance atoms other than O and H



- ◆ **Step (5) :** Balance O atoms



- ◆ **Step (6) :** Balance H atoms as is done in acidic medium.



- (b) **Ion electron method or half cell method :**

By this method redox equation are balanced in two different medium


- (a) Acidic medium (b) Basic medium

- ◆ **Balancing in acidic medium**


Students are advised to follow the following steps to balance the redox reactions by ion electron method in acidic medium.

### □ Solved Examples :


**Ex :** Balance the following redox reaction.




**Sol : Step-I** assign the oxidation No. to each elements present in the reaction.



**Step-II** Now convert the reaction in ionic form by eliminating the elements or species which are not going either oxidation or reduction



**Step-III** Now identify the oxidation / reduction occurring into the reaction.



**Step-IV** Split the ionic reaction in two half one for oxidation and other for reduction



**Step-V** Balance the atom other than oxygen and hydrogen atom in both half reactions





Fe & Mn atom are balanced in both side.

**Step-VI** Now balance O & H atom by  $\text{H}_2\text{O}$  &  $\text{H}^+$  respectively by the following way for one excess oxygen atom add one  $\text{H}_2\text{O}$  on the other side and two  $\text{H}^+$  on the same side.





**Step VII** Now see equation (i) & (ii) is balanced atomwise. Now balance both equations chargewise to balance the charge add electrons to the electrically positive side.



**Step VIII** The number of electrons gained and lost in each half-reaction are equalised by multiply suitable factor in both the half reaction and finally the half reactions are added to give the over all balanced reaction.



Here we multiply equation (i) by 5 and (ii) by one



(Here at this stage you will get balanced redox reaction in ionic form)


**Step IX** Now convert the ionic reaction in to molecular form by adding the elements or species which are removed in step (II).

Now by some manipulation you will get



◆ **Balancing in basic medium :**

In this case except step VI all the steps are same. We can understand it by following example balance the redox reaction in basic medium



**Sol :** By using up to step V we will get

- 1.**      (i) 1                      (ii) 2                      (iii) 1

◆ **For Base :**

v.f. = number of replicable  $\text{OH}^-$  ions

□ **Solved Examples :**

**Ex :**  $\text{NaOH}$ ,  $\text{KOH}$

**Sol :** v.f.  $\rightarrow$  1 1

E.  $\rightarrow$   $\frac{M}{1}$   $\frac{M}{1}$

**Self practice problems :**

1. Find the valence factor for following bases

(i)  $\text{Ca(OH)}_2$

(ii)  $\text{CsOH}$

(iii)  $\text{Al(OH)}_3$

**Answers :**

1. (i) 2 (ii) 1 (iii) 3

◆ **Acid - base reaction :**

In case of acid base reaction, the valence factor is the actual number of  $\text{H}^+$  or  $\text{OH}^-$  replaced in the reaction. The acid or base may contain more number of replaceable  $\text{H}^+$  or  $\text{OH}^-$  than actually replaced in reaction.

v.f. for base is the number of  $\text{H}^+$  ion from the acid replaced by per molecule of the base.

□ **Solved Examples :**

**Ex :**  $2\text{NaOH} + \text{H}_2\text{SO}_4 \rightarrow \text{Na}_2\text{SO}_4 + 2\text{H}_2\text{O}$   
Base acid

**Sol :** valency factor of base = 1

valency factor of acid = 2

Here two molecule of  $\text{NaOH}$  replaced  $2\text{H}^+$  ion from the  $\text{H}_2\text{SO}_4$  therefore per molecule of  $\text{NaOH}$  replaced only one  $\text{H}^+$  ion of acid so v.f. = 1

◆ v.f. for acid is number of  $\text{OH}^-$  replaced for the base by per molecule of acid

**Ex :**  $\text{NaOH} + \text{H}_2\text{SO}_4 \rightarrow \text{NaHSO}_4 + \text{H}_2\text{O}$   
Base acid

**Sol :** valence factor of acid = 1

here one of molecule of  $\text{H}_2\text{SO}_4$  replaced one  $\text{OH}^-$  from  $\text{NaOH}$  therefore v.f. for  $\text{H}_2\text{SO}_4$  is = 1

$$E = \frac{\text{mol. wt. of } \text{H}_2\text{SO}_4}{1}$$

◆ **For Salts :**

(a) In non reacting condition

v.f. = Total number of positive charge or negative charge present into the compound.

□ **Solved Examples :**


**Ex :**  $\text{Na}_2\text{CO}_3$   $\text{Fe}_2(\text{SO}_4)_3 (2\text{Fe}^{3+} + 3\text{SO}_4^{2-})$   $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$

**Sol :** V.f. 2 2 3 = 6 2

E.  $\frac{M}{2}$   $\frac{M}{6}$   $\frac{M}{2}$

(b) Salt in reacting condition :

❑ **Solved Examples :**



**Sol :** It is a acid base reaction therefore v.f. for  $\text{Na}_2\text{CO}_3$  is one while in non reaction condition it will be two.


(c) Eq. wt. of oxidising / reducing agents in redox reaction :

The equivalent weight of an oxidising agent is that weight which accepts one mole electron in a chemical reaction.

(a) Equivalent wt. of an oxidant (get reduced)

$$= \frac{\text{Mol.wt.}}{\text{No. of electrons gained by one mole}}$$

**Ex :** In acidic medium



$$\text{Eq. wt. of K}_2\text{Cr}_2\text{O}_7 = \frac{\text{Mol. wt of K}_2\text{Cr}_2\text{O}_7}{6} = \frac{\text{Mol. wt.}}{6}$$

**Note :** [6 in denominator indicates that 6 electrons were gained by  $\text{Cr}_2\text{O}_7^{2-}$  as it is clear from the given balanced equation]

(b) Similarly equivalent wt. of a reductant (gets oxidised)

$$= \frac{\text{Mol. wt.}}{\text{No. of electrons lost by one mole}}$$

**Ex:** In acidic medium,  $\text{C}_2\text{O}_4^{2-} \longrightarrow 2\text{CO}_2 + 2e^-$

$$\text{C}_2\text{O}_4^{2-} \longrightarrow 2\text{CO}_2 + 2e^-$$

Here, Total electrons lost = 2

$$\text{So, eq. wt.} = \frac{\text{Mol. wt.}}{2}$$

(c) In different condition a compound may have different equivalent weights. Because, it depends upon the number of electrons gained or lost by that compound in that reaction.

**Ex :** (i)  $\text{MnO}_4^- \longrightarrow \text{Mn}^{2+}$  (acidic medium)

(+7)                      (+2)

Here 5 electrons are taken so eq. wt. =  $\frac{\text{Mol.wt.of KMnO}_4}{5} = \frac{158}{5} = 31.6$

(ii)  $\text{MnO}_4^- \longrightarrow \text{Mn}^{+4}$  (neutral medium)  
(+7) (+4)

Here, only 3 electrons are gained, so eq. wt. =  $\frac{\text{Mol.wt. of KMnO}_4}{3} = \frac{158}{3} = 52.7$

(iii)  $\text{MnO}_4^- \longrightarrow \text{MnO}_4^{2-}$  (alkaline medium)

(+7)                      (+6)

Here, only one electron is gained, so eq. wt. =  $\frac{\text{Mol.wt. of KMnO}_4}{1} = 158$

**Note :** It is important to note that  $\text{KMnO}_4$  acts as an oxidant in every medium although with different strength which follows the order.

acidic medium > neutral medium > alkaline medium

**Ex :** 
$$2\text{S}_2\text{O}_3^{--} \longrightarrow \text{S}_4\text{O}_6^{--} + 2\text{e}^-$$
  
(Reducing agent)

$$\text{equivalent weight of } \text{S}_2\text{O}_3^{--} = \frac{2M}{2} = M$$

◆ **Questions based on Equivalent weight :**

- Molecular weight of  $\text{KMnO}_4$  in acidic medium and neutral medium will be respectively :  
 (A) 7 equivalent wt. and 2 equivalent wt.  
 (B) 5 equivalent wt. and 4 equivalent wt.  
 (C) 4 equivalent wt. and 5 equivalent wt.  
 (D) 2 equivalent wt. and 4 equivalent wt.
- Equivalent wt. of  $\text{H}_3\text{PO}_4$  in each of the reaction will be respectively :  
 $\text{H}_3\text{PO}_4 + \text{OH}^- \longrightarrow \text{H}_2\text{PO}_4^- + \text{H}_2\text{O}$   
 $\text{H}_3\text{PO}_4 + 2\text{OH}^- \longrightarrow \text{HPO}_4^{2-} + 2\text{H}_2\text{O}$   
 $\text{H}_3\text{PO}_4 + 3\text{OH}^- \longrightarrow \text{PO}_4^{3-} + 3\text{H}_2\text{O}$   
 (A) 98, 49, 32.67      (B) 49, 98, 32.67      (C) 98, 32.67, 49      (D) 32.67, 49, 98
- In acidic medium, equivalent weight of  $\text{K}_2\text{Cr}_2\text{O}_7$  (Mol. wt. = M) is –  
 (A)  $M/3$       (B)  $M/4$       (C)  $M/6$       (D)  $M/2$

**Answers :**

- (1) B      (2) A      (3) C

□ **NORMALITY :**

Normality of solution is defined as the number of equivalent of solute present in one litre (1000 mL) solutions.  
 Let a solution is prepared by dissolving W g of solute of eq. wt. E in V mL water.

- ◆ No. of equivalent of solute =  $\frac{W}{E}$
- ◆ V mL of solution have  $\frac{W}{E}$  equivalent of solute
- ◆ 1000 mL solution have  $\frac{W \times 1000}{E \times V \text{ mL}}$
- ◆ **Normality (N) =  $\frac{W \times 1000}{E \times V \text{ mL}}$**
- ◆ **Normality (N) = Molarity    Valence factor**  
 $\text{Normality (N)} = \text{molarity} \times \text{Valence factor (n)}$   
 or  $N \times V \text{ (in mL)} = M \times V \text{ (in mL)} \times n$   
 or milli equivalents = millimoles  $\times n$

□ **Solved Examples :**

**Ex :** Calculate the normality of a solution containing 15.8 g of  $\text{KMnO}_4$  in 50 mL acidic solution.

**Sol :** **Normality (N) =  $\frac{W \times 1000}{E \times V \text{ mL}}$**

where  $W = 15.8 \text{ g}$ ,       $V = 50 \text{ mL}$        $E = \frac{\text{molar mass of } \text{KMnO}_4}{\text{Valence factor}} = 158/5 = 31.6$

So,  $N = 10$

**Ex :** Calculate the normality of a solution containing 50 mL of 5 M solution  $\text{K}_2\text{Cr}_2\text{O}_7$  in acidic medium.

**Sol :** **Normality (N) = Molarity    Valence factor**

$= 5 \times 6 = 30 \text{ N}$


❑ **MOLARITY V/S NORMALITY :**

Molarity (M)		Normality (N)	
1.	No. of moles of solute present in one litre of solution.	1.	No. of equivalents of solute present in one litre of solution.
2.	No. of moles = $\frac{W}{M}$	2.	No. of equivalents = $\frac{W}{E}$
3.	$\frac{W}{M} \times 1000 =$ No. of millimoles	3.	$\frac{W}{E} \times 1000 =$ No. of equivalence
4.	Molarity $V(\text{in mL}) =$ No. of millimoles	4.	Normality $V(\text{in mL}) =$ No. of equivalents
5.	Molarity = $\frac{\text{milli moles}}{\text{Volume of solution in mL}}$	5.	Normality = $\frac{\text{milli equivalents}}{\text{Volume of solution in mL}}$

❑ **LAW OF EQUIVALENCE :**

The law states that one equivalent of an element combine with one equivalent of the other, and in a chemical reaction equivalent and mill equivalent of reactants react in equal amount to give same no. of equivalent or milli equivalents of products separately.

**According :**




$$m.\text{eq of A} = m.\text{eq of B} = m.\text{eq of M} = m.\text{eq of N}$$



$$m.\text{eq of } M_xN_y = m.\text{eq of M} = m.\text{eq of N}$$

❑ **Solved Examples :**

**Ex :** Find the number of moles of  $\text{KMnO}_4$  needed to oxidise one mole  $\text{Cu}_2\text{S}$  in acidic medium.



**Sol :** From law of equivalence

$$\text{equivalents of } \text{Cu}_2\text{S} = \text{equivalents of } \text{KMnO}_4$$

$$\text{moles of } \text{Cu}_2\text{S} \times \text{v.f} = \text{moles of } \text{KMnO}_4 \times \text{v.f.}$$

$$\text{moles of } \text{Cu}_2\text{S} \times 8 = 1 \times 5 \Rightarrow \text{moles of } \text{Cu}_2\text{S} = 5/8$$

**Ex :** The number of moles of oxalate ions oxidized by one mole of  $\text{MnO}_4^-$  ion in acidic medium.

(A)  $\frac{5}{2}$

(B)  $\frac{2}{5}$

(C)  $\frac{3}{5}$

(D)  $\frac{5}{3}$

**Sol :** Equivalents of  $\text{C}_2\text{O}_4^{2-} =$  equivalents of  $\text{MnO}_4^-$

$$x (\text{mole}) \times 2 = 1 \times 5$$

$$x = \frac{5}{2}$$

**Ex.** What volume of 6 M HCl and 2 M HCl should be mixed to get two litre of 3 M HCl ?

**Sol.** Let, the volume of 6 M HCl required to obtain 2 L of 3M HCl = x L

$$\therefore \text{Volume of 2 M HCl required} = (2 - x) \text{ L}$$

$$\begin{array}{ccccc} M_1V_1 & + & M_2V_2 & = & M_3V_3 \\ 6M \text{ HCl} & & 2M \text{ HCl} & & 3M \text{ HCl} \end{array}$$

$$6(x) + 2(2 - x) = 3 \times 2$$

$$\Rightarrow 6x + 4 - 2x = 6 \Rightarrow 4x = 2$$

$$\therefore x = 0.5 \text{ L}$$

Hence, volume of 6 M HCl required = 0.5 L

Volume of 2M HCl required = 1.5 L



**Ex.** In a reaction vessel, 1.184 g of NaOH is required to be added for completing the reaction. How many millilitre of 0.15 M NaOH should be added for this requirement ?

**Sol.** Amount of NaOH present in 1000 mL of 0.15 M NaOH =  $0.15 \times 40 = 6$  g

$$\therefore 1 \text{ mL of this solution contain NaOH} = \frac{6}{1000} \times 10^{-3} \text{ g}$$

$$\therefore 1.184 \text{ g of NaOH will be present in} = \frac{1}{6 \times 10^{-3}} \times 1.184 = 197.33 \text{ mL}$$

**Ex.** What weight of  $\text{Na}_2\text{CO}_3$  of 85% purity would be required to prepare 45.6 mL of 0.235N  $\text{H}_2\text{SO}_4$  ?

**Sol.** Meq. of  $\text{Na}_2\text{CO}_3$  = Meq. of  $\text{H}_2\text{SO}_4$

$$= 45.6 \times 0.235$$

$$\therefore \frac{W_{\text{Na}_2\text{CO}_3}}{E_{\text{Na}_2\text{CO}_3}} \times 1000 = 45.6 \times 0.235$$

$$\Rightarrow \frac{W_{\text{Na}_2\text{CO}_3}}{106/2} \times 1000 = 45.6 \times 0.235$$

$$\therefore W_{\text{Na}_2\text{CO}_3} = 0.5679 \text{ g}$$


For 85 g of pure  $\text{Na}_2\text{CO}_3$ , weighed sample = 100 g

$$\therefore \text{For } 0.5679 \text{ g of pure } \text{Na}_2\text{CO}_3, \text{ weighed sample} = \frac{100}{85} \times 0.5679$$


$$= 0.6681 \text{ g}$$

#### ❑ Drawbacks of Equivalent concept :

- ◆ Since equivalent weight of a substance for example oxidising or reducing agent may be variable hence it is better to use mole concept.



$$\therefore \text{Eq. wt of } \text{MnO}_4^- = \frac{\text{MnO}_4^-(\text{mol.wt.})}{5}$$



$$\therefore \text{Eq. wt. of } \text{MnO}_4^- = \frac{\text{MnO}_4^-}{3}$$



Thus the no. of equivalents of  $\text{MnO}_4^-$  will be different in the above two cases but no. of moles will be same.

- ◆ Normality of any solution depends on reaction while molarity does not.

For example.

Consider 0.1 mol  $\text{KMnO}_4$  dissolved in water to make 1L solution. Molarity of this solution is 0.1 M. However, its normality is not fixed. It will depend upon the reaction in which  $\text{KMnO}_4$  participates, e.g. If  $\text{KMnO}_4$  forms  $\text{Mn}^{2+}$ , normality =  $0.1 \times 5 = 0.5$  N. This same sample of  $\text{KMnO}_4$ , if employed in a reaction giving  $\text{MnO}_2$  as product (Mn in +4 state) will have normality  $0.1 \times 3 = 0.3$  N.

- ◆ The concept of equivalents is handy, but it should be used with care. One must never equate equivalents in a sequence which involves same element in more than two oxidation states. Consider an example  $\text{KIO}_3$  reacts with KI to liberate iodine and liberated iodine is titrated with standard hypo solution, The reaction are



meq of hypo = meq of  $\text{I}_2$  = meq of  $\text{IO}_3^-$  + meq of  $\text{I}^-$

$\therefore \text{IO}_3^-$  react with  $\text{I}^- \Rightarrow$  meq of  $\text{IO}_3^-$  = meq of  $\text{I}^-$

◆ meq of hypo = 2 meq of  $\text{IO}_3^-$

This is wrong. Note that  $\text{I}_2$  formed by (i) have v.f. = 5/3 & reacted in equation (ii) have v.f. = 2.

$\therefore$  v.f. of  $\text{I}_2$  in both the equation are different therefore we cannot equate m.eq is sequence. In this type of case students are advised to use mole concept.

#### □ Solved Example :

**Ex :** How many milliliters of 0.02000 M  $\text{KMnO}_4$  solution would be required to exactly titrate 25.00 mL of 0.2000 M  $\text{Fe}(\text{NO}_3)_2$  solution.

**Sol : Method - 1 : Mole concept method**


Starting with 25.00 mL of 0.2000M  $\text{Fe}^{2+}$ , we can write.

Millimoles of  $\text{Fe}^{2+}$  = 25.00 0.2000

and in volume V (in milliliters of the  $\text{MnO}_4^-$ )

Millimoles of  $\text{MnO}_4^-$  = V (0.02000)

The balanced reaction is :



This requires that at the equivalent point,

$$\frac{V(0.02000)}{1} = \frac{(25.00)(0.2000)}{5} \quad V = 50.00 \text{ mL}$$

**Method - 2 : Equivalent Method :**

Equivalents of  $\text{MnO}_4^-$  = 5 moles of  $\text{MnO}_4^-$

Normality  $\text{MnO}_4^-$  = 5 molarity of  $\text{MnO}_4^-$

For  $\text{Fe}^{2+}$ , moles and equivalents are equal,

At the equivalence point,

Equivalents of  $\text{MnO}_4^-$  = Equivalents of  $\text{Fe}^{2+}$

or  $V_{\text{MnO}_4^-}$  Normality of  $\text{MnO}_4^-$

=  $V_{\text{Fe}^{2+}}$  Normality of  $\text{Fe}^{2+}$

For 0.02000 M  $\text{MnO}_4^-$  solution


Normality of  $\text{MnO}_4^-$  = (5) (0.02000) = 0.1 N and for 0.2000 M  $\text{Fe}^{2+}$  solution

Normality of  $\text{Fe}^{2+}$  = 0.2000 N

$$V_{\text{MnO}_4^-} = (25.00 \text{ mL}) \left( \frac{0.2000}{0.1000} \right) = 50.00 \text{ mL}$$


#### □ HYDROGEN PEROXIDE ( $\text{H}_2\text{O}_2$ ) :

$\text{H}_2\text{O}_2$  can behave both like oxidising and reducing agents in both the medium (acidic and basic).




◆ **Oxidising agent : ( $\text{H}_2\text{O}_2$ )**

(a) **Acidic medium :**



$$\text{v.f.} = 2$$


(b) **Basic medium :**



$$\text{v.f.} = 2$$


◆ **Reducing agent : ( $\text{H}_2\text{O}_2$ ) :**

(a) **Acidic medium :**



$$\text{v.f.} = 2$$

(b) **Basic medium :**



$$\text{v.f.} = 2$$

**Note :** Valency factor of  $\text{H}_2\text{O}_2$  is always equal to 2.

**Volume strength of  $\text{H}_2\text{O}_2$  :** Strength of  $\text{H}_2\text{O}_2$  is represented as 10 V, 20 V, 30 V etc.

**20V  $\text{H}_2\text{O}_2$**  means **one litre** of this sample of  $\text{H}_2\text{O}_2$  on decomposition gives **20 L of  $\text{O}_2$**  gas at **S.T.P.**

$$\text{Normality of } \text{H}_2\text{O}_2 \text{ (N)} = \frac{\text{Volume, strength of } \text{H}_2\text{O}_2}{5.6}$$

$$\therefore M_{\text{H}_2\text{O}_2} = \frac{N_{\text{H}_2\text{O}_2}}{\text{v.f.}} = \frac{N_{\text{H}_2\text{O}_2}}{2}$$

$$\text{◆ Molarity of } \text{H}_2\text{O}_2 \text{ (M)} = \frac{\text{Volume, strength of } \text{H}_2\text{O}_2}{11.2}$$

**Strength (in g/L) :** Denoted by S

Strength = molarity mol. wt.

$$= \text{molarity} \times 34$$

Strength = Normality Eq. weight.

$$= \text{Normality} \times 17$$

□ **Solved Example :**

**Ex :** 20 mL of  $\text{H}_2\text{O}_2$  after acidification with dil  $\text{H}_2\text{SO}_4$  required 30 mL of  $\frac{\text{N}}{12}$   $\text{KMnO}_4$  for complete oxidation. The strength of  $\text{H}_2\text{O}_2$  solution is [Molar mass of  $\text{H}_2\text{O}_2 = 34$ ]

$$\text{Sol :} \quad 30 \times \frac{1}{12} = 20 \times \text{N}'$$


$$\text{N}' = \frac{30}{12 \times 20} = \frac{1}{8}$$

$$\therefore \text{strength} = \text{N}' \times \text{equivalent mass} = \frac{1}{8} \times 17 = 2.12 \text{ g/L}$$

**Hardness of water (Hard water does not give lather with soap)**

Temporary hardness - due to bicarbonates of Ca & Mg

Permanent hardness - due to chloride & sulphates of Ca & Mg. There are some method by which we can softening the water



#### □ PARTS PER MILLION (ppm) :

When the solute is present in very less amount then this concentration term is used. It is defined as the number of parts of the solute present in every 1 million parts of the solution. ppm can both be in terms of mass or in terms of moles. If nothing has been specified we take ppm to be in terms of mass. hence a 100 ppm solution means that 100 g of solute are present in every 1000000 g of solution.

$$\text{ppm}_A = \frac{\text{mass of A}}{\text{Total mass}} \quad 10^6 = \text{mass fraction} \quad 10^6$$

#### ◆ Measurement of Hardness :

Hardness is measured in terms of ppm (parts per million) of  $\text{CaCO}_3$  or equivalent to it.

#### □ Solved Example :

**Ex :** 0.00012%  $\text{MgSO}_4$  and 0.000111%  $\text{CaCl}_2$  is present in water. What is the measured hardness of water and millimoles of washing soda requires to purify 1000 litre water.

**Sol :** 0.00012%  $\text{MgSO}_4 = 1 \text{ mg } \text{CaCO}_3 \text{ 1 L water}$

0.000111%  $\text{CaCl}_2 = 1 \text{ mg } \text{CaCO}_3 \text{ 1 L water}$





hardness = 2 ppm and mm of  $\text{Na}_2\text{CO}_3$  require is 20

#### ◆ Strength of Oleum


Oleum is  $\text{SO}_3$  dissolved in 100%  $\text{H}_2\text{SO}_4$ . Sometimes, oleum is reported as more than 100% by weight, say y% (where  $y > 100$ ). This means that  $(y - 100)$  grams of water, when added to 100 g of given oleum sample, will combine with all the free  $\text{SO}_3$  in the oleum to give 100% sulphuric acid. Hence weight % of free  $\text{SO}_3$  in oleum =  $80(y - 100)/18$

#### □ CALCULATION OF AVAILABLE CHLORINE FROM A SAMPLE OF BLEACHING POWDER :

The weight % of available  $\text{Cl}_2$  from the given sample of bleaching powder on reaction with dil acids or  $\text{CO}_2$  is called available chlorine.



#### ◆ Method of determination :



$$\% \text{ of } \text{Cl}_2 = \frac{3.55 \times x \times V(\text{mL})}{W(\text{g})}$$

where  $x$  = molarity of hypo solution

$v$  = mL of hypo solution used in titration.

#### □ Solved Example :

**Ex :** 3.55 g sample of bleaching powder suspended in  $\text{H}_2\text{O}$  was treated with enough acetic acid and KI solution.

Iodine thus liberated requires 80 mL of 0.2 M hypo for titration. Calculate the % of available chlorine.

[Available Chlorine = mass of chlorine liberated / mass of bleaching powder 100]

**Sol :** moles of iodine = moles of chlorine =  $\frac{80 \times 0.2}{2} \times 10^{-3} = 8 \times 10^{-3}$

$$\text{so required \%} = \frac{8 \times 71 \times 10^{-3}}{3.55} \times 100\% = 16\%$$

❑ **FOR ACID-BASE (NEUTRALIZATION REACTION) OR REDOX REACTION :**

$N_1V_1 = N_2V_2$  is always true.

But  $M_1V_1 = M_2V_2$  (may or may not be true )

But  $M_1 n_1 V_1 = M_2 n_2 V_2$  (always true where n terms represent n-factor).

❑ **'n' FACTOR : FACTOR RELATING MOLECULAR WEIGHT AND EQUIVALENT WEIGHT :**

$$\text{n-factor} = \frac{M}{E}$$

$$E = \frac{M}{\text{n-factor}}$$

❑ **n-FACTOR IN VARIOUS CASES**

**In Non Redox Change**

◆ **n-factor for element** : Valency of the element

◆ **For acids** : Acids will be treated as species which furnish  $H^+$  ions when dissolved in a solvent. The n-factor of an acid is the no. of acidic  $H^+$  ions that a molecule of the acid would give when dissolved in a solvent (Basicity).

For example, for HCl (n = 1),  $HNO_3$  (n = 1),  $H_2SO_4$  (n = 2),  $H_3PO_4$  (n = 3) and  $H_3PO_3$  (n = 2)

◆ **For bases** : Bases will be treated as species which furnish  $OH^-$  ions when dissolved in a solvent . The n-factor of a base is the no. of  $OH^-$  ions that a molecule of the base would give when dissolved in a solvent (**Acidity**).

For example, NaOH (n = 1),  $Ba(OH)_2$  (n = 2),  $Al(OH)_3$  (n = 3), etc.

◆ **For salts** : A salt reacting such that no atom of the salt undergoes any change in oxidation state.

For example,  $2AgNO_3 + MgCl_2 \rightarrow Mg(NO_3)_2 + 2AgCl$

In this reaction, it can be seen that the oxidation state of Ag, N, O, Mg and Cl remains the same even in the product. The n-factor for such a salt is the total **charge on cation or anion**.

**In Redox Change**

For oxidizing agent or reducing agent n-factor is the **change in oxidation number per mole of the substance**.

❑ **SOME OXIDIZING AGENTS/REDUCING AGENTS WITH EQ. WT.**

Species	Changed to	Reaction	Electrons exchanged or change in O.N.	Eq. wt.
$MnO_4^-$ (O.A.)	$Mn^{+2}$ in acidic medium	$MnO_4^- + 8H^+ + 5e^- \longrightarrow Mn^{2+} + 4H_2O$	5	$E = \frac{M}{5}$
$MnO_4^-$ (O.A.)	$MnO_2$ in neutral medium	$MnO_4^- + 3e^- + 2H_2O \longrightarrow MnO_2 + 4OH^-$	3	$E = \frac{M}{3}$
$MnO_4^-$ (O.A.)	$MnO_4^{2-}$ in basic medium	$MnO_4^- + e^- \longrightarrow MnO_4^{2-}$	1	$E = \frac{M}{1}$
$Cr_2O_7^{2-}$ (O.A.)	$Cr^{3+}$ in acidic medium	$CrO_7^{2-} + 14H^+ + 6e^- \longrightarrow 2Cr^{3+} + 7H_2O$	6	$E = \frac{M}{6}$
$MnO_2$ (O.A.)	$Mn^{2+}$ in acidic medium	$MnO_2 + 4H^+ + 2e^- \longrightarrow Mn^{2+} + 2H_2O$	2	$E = \frac{M}{2}$

$\text{Cl}_2$ (O.A.) in bleaching powder	$\text{Cl}^-$	$\text{Cl}_2 + 2e^- \longrightarrow 2\text{Cl}^-$	2	$E = \frac{M}{2}$
$\text{CuSO}_4$ (O.A.) in iodometric titration	$\text{Cu}^+$	$\text{Cu}^{2+} + e^- \longrightarrow \text{Cu}^+$	1	$E = \frac{M}{1}$
$\text{S}_2\text{O}_3^{2-}$ (R.A.)	$\text{S}_4\text{O}_6^{2-}$	$2\text{S}_2\text{O}_3^{2-} \longrightarrow \text{S}_4\text{O}_6^{2-} + 2e^-$ (for two molecules)	2	$E = \frac{2M}{2} = M$
$\text{H}_2\text{O}_2$ (O.A.)	$\text{H}_2\text{O}$	$\text{H}_2\text{O}_2 + 2\text{H}^+ + 2e^- \longrightarrow 2\text{H}_2\text{O}$	2	$E = \frac{M}{2}$
$\text{H}_2\text{O}_2$ (R.A.)	$\text{O}_2$	$\text{H}_2\text{O}_2 \longrightarrow \text{O}_2 + 2\text{H}^+ + 2e^-$ (O.N. of oxygen in $\text{H}_2\text{O}_2$ is -1 per atom)	2	$E = \frac{M}{2}$
$\text{Fe}^{2+}$ (R.A.)	$\text{Fe}^{3+}$	$\text{Fe}^{2+} \longrightarrow \text{Fe}^{3+} + e^-$	1	$E = \frac{M}{1}$

**Ex.** To find the n-factor in the following chemical changes.

- (i)  $\text{KMnO}_4 \xrightarrow{\text{H}^+} \text{Mn}^{2+}$       (ii)  $\text{KMnO}_4 \xrightarrow{\text{H}_2\text{O}} \text{Mn}^{4+}$   
 (iii)  $\text{KMnO}_4 \xrightarrow{\text{OH}^-} \text{Mn}^{6+}$       (iv)  $\text{K}_2\text{Cr}_2\text{O}_7 \xrightarrow{\text{H}^+} \text{Cr}^{3+}$   
 (v)  $\text{C}_2\text{O}_4^{2-} \rightarrow \text{CO}_2$       (vi)  $\text{FeSO}_4 \rightarrow \text{Fe}_2\text{O}_3$   
 (vii)  $\text{Fe}_2\text{O}_3 \rightarrow \text{FeSO}_4$

**Sol.** (i) In this reaction,  $\text{KMnO}_4$  which is an oxidizing agent, itself gets reduced to  $\text{Mn}^{2+}$  under acidic conditions.

$$n = |1 \quad (+7) - 1 \quad (+2)| = 5$$

(ii) In this reaction,  $\text{KMnO}_4$  gets reduced to  $\text{Mn}^{4+}$  under neutral or slightly (weakly) basic conditions.

$$n = |1 \quad (+7) - 1 \quad (+4)| = 3$$

(iii) In this reaction,  $\text{KMnO}_4$  gets reduced to  $\text{Mn}^{6+}$  under basic conditions.

$$n = |1 \quad (+7) - 1 \quad (+6)| = 1$$

(iv) In this reaction,  $\text{K}_2\text{Cr}_2\text{O}_7$  which acts as an oxidizing agent reduced to  $\text{Cr}^{3+}$  under acidic conditions. (It does not react under basic conditions.)

$$n = |2 \quad (+6) - 2 \quad (+3)| = 6$$

(v) In this reaction,  $\text{C}_2\text{O}_4^{2-}$  (oxalate ion) gets oxidized to  $\text{CO}_2$  when it is reacted with an oxidizing agent.

$$n = |2 \quad (+3) - 2 \quad (+4)| = 2$$

(vi) In this reaction, ferrous ions get oxidized to ferric ions.

$$n = |1 \quad (+2) - 1 \quad (+3)| = 1$$

(vi) In this reaction, ferric ions are getting reduced to ferrous ions.

$$n = |2 \quad (+3) - 2 \quad (+2)| = 2$$

**Ex.** Calculate the molar ratio in which the following two substances would react ?

$\text{Ba}_3(\text{PO}_4)_2$  and  $\text{AlCl}_3$

**Sol.** n-factor of  $\text{Ba}_3(\text{PO}_4)_2 = 3 \quad (+2) = 6 = n_1$

While n-factor of  $\text{AlCl}_3 = 1 \quad (+3) = 3 = n_2$

$$\frac{n_1}{n_2} = \frac{6}{3} \quad \text{If} \quad \frac{n_1}{n_2} = \frac{x}{y}$$

Molar ratio  $= \frac{y}{x}$  (inverse of equivalent ratio)

$\therefore$  Molar ratio in which  $\text{Ba}_3(\text{PO}_4)_2$  and  $\text{AlCl}_3$  will react  $= 3 : 6 = 1 : 2$

## □ APPLICATIONS OF THE LAW OF EQUIVALENCE

### Simple Titration

In this, we can find the concentration of unknown solution by reacting it with solution of known concentration (standard solution).

For example, let there be a solution of substance A of unknown concentration. We are given solution of another substance B whose concentration is known ( $N_1$ ). We take a certain known volume of A in a flask ( $V_2$ ) and then we add B to A slowly till all the A is consumed by B (this can be known with the help of indicators). Let us, assume that the volume of B consumed is  $V_1$ . According to the law of equivalence, the number of g equivalents of B at the end point.

$$\therefore N_1 V_1 = N_2 V_2, \text{ where } N_2 \text{ is the conc. of A.}$$

From this we can calculate the value of  $N_2$ .

**Ex.** 0.4 M  $\text{KMnO}_4$  solution completely reacts with 0.05 M  $\text{FeSO}_4$  solution under acidic conditions. The volume of  $\text{FeSO}_4$  used is 50 mL. What volume of  $\text{KMnO}_4$  was used ?

**Sol.** 0.4      5       $V = 0.05$       50

$$V = 1.25 \text{ mL}$$

**Ex.** 1.20 g sample of  $\text{Na}_2\text{CO}_3$  and  $\text{K}_2\text{CO}_3$  was dissolved in water to form 100 mL of a Solution. 20 mL of this solution required 40 mL of 0.1 N HCl for complete neutralization. Calculate the weight of  $\text{Na}_2\text{CO}_3$  in the mixture. If another 20 mL of this solution is treated with excess of  $\text{BaCl}_2$  what will be the weight of the precipitate ?

**Sol.** Let, weight of  $\text{Na}_2\text{CO}_3 = x$  g

Weight of  $\text{K}_2\text{CO}_3 = y$  g

$$\therefore x + y = 1.20 \text{ g} \quad \dots\dots\dots(i)$$

For neutralization reaction of 100 mL

Meq. of  $\text{Na}_2\text{CO}_3 + \text{Meq. of } \text{K}_2\text{CO}_3 = \text{Meq. of HCl}$

$$\Rightarrow \frac{x}{106} \times 2 \times 1000 + \frac{y}{138} \times 2 \times 1000 = \frac{40 \times 0.1 \times 100}{20}$$

$$\therefore 69x + 53y = 73.14 \quad \dots\dots\dots(ii)$$

From Eqs. (i) and (ii), we get

$$x = 0.5962 \text{ g}$$

$$y = 0.604 \text{ g}$$

Solution of  $\text{Na}_2\text{CO}_3$  and  $\text{K}_2\text{CO}_3$  gives ppt. of  $\text{BaCO}_3$  with  $\text{BaCl}_2$

(Meq. of  $\text{Na}_2\text{CO}_3 + \text{Meq. of } \text{K}_2\text{CO}_3$ ) in 20 mL = Meq. of  $\text{BaCO}_3$

$\Rightarrow$  Meq. of HCl for 20 mL mixture = Meq. of  $\text{BaCO}_3$

$$\Rightarrow \text{Meq. of } \text{BaCO}_3 = 40 \times 0.1 = 4$$

$$\frac{W_{\text{BaCO}_3}}{M_{\text{BaCO}_3}} \times 1000 = 4 \times 100 = 400$$

$$\frac{W_{\text{BaCO}_3}}{197} \times 2 \times 1000 = 400$$

$$\therefore W_{\text{BaCO}_3} = 0.394 \text{ g}$$

## ❑ BACK TITRATION

Back titration is used to calculate % purity of a sample. Let us assume that we are given an impure solid substance C weighing  $w$  gs and we are asked to calculate the percentage of pure C in the sample. We will assume that the impurities are inert. We are provided with two solutions A and B, where the concentration of B is known ( $N_1$ ) and that of A is not known. This type of titration will work only if the following condition is satisfied, i.e. the nature of compounds A, B and C should be such that A and B can react with each other. A and C can react with each other but the product of A and C should not react with B.

Now, we take a certain volume of A in a flask (the g equivalents of A taken should be  $\geq$  g equivalents of C in the sample and this can be done by taking A in excess). Now, we perform a simple titration using B. Let us assume that the volume of B used is  $V_1$ . In another beaker, we again take the solution of A in the same volume as taken earlier. Now, C is added to this and after the reaction is completed, the solution is being titrated with B. Let us assume that the volume of B used up is  $V_2$ . Gram equivalents of B used in the first titration =  $N_1 V_1$ .

$$\therefore \text{gm. equivalents of A initially} = N_1 V_1$$

$$\text{gm. equivalents of B used in the second titration} = N_1 V_2$$

$$\therefore \text{gm. equivalents of A left in excess after reacting with C} = N_1 V_2$$

$$\text{gm. equivalents of A that reacted with C} = N_1 V_1 - N_1 V_2$$

$$\text{If the n-factor of C is } x, \text{ then the moles of pure C} = \frac{N_1 V_1 - N_1 V_2}{x}$$

$$\therefore \text{The weight of C} = \frac{N_1 V_1 - N_1 V_2}{x} \quad \text{Molecular weight of C}$$

$$\therefore \text{Percentage of C} = \frac{N_1 V_1 - N_1 V_2}{x} \times \frac{\text{Molecular wt. of C}}{w} \times 100$$

## ❑ SOME REDOX TITRATIONS (EXCLUDING IODOMETRIC / IODIMETRIC)

Estimation of	By titrating with	Reactions	Relation *between O.A. and R.A.
$\text{Fe}^{2+}$	$\text{MnO}_4^-$	$\text{Fe}^{2+} \longrightarrow \text{Fe}^{3+} + e^-$ $\text{MnO}_4^- + 8\text{H}^+ + 5e^- \longrightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$	$5\text{Fe}^{2+} \equiv \text{MnO}_4^-$ Eq. wt. $\text{Fe}^{2+} = M/1$ Eq. wt. $\text{MnO}_4^- = M/5$
$\text{Fe}^{2+}$	$\text{Cr}_2\text{O}_7^{2-}$	$\text{Fe}^{2+} \longrightarrow \text{Fe}^{3+} + e^-$ $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6e^- \longrightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$	$6\text{Fe}^{2+} \equiv \text{Cr}_2\text{O}_7^{2-}$ Eq. wt. $\text{Cr}_2\text{O}_7^{2-} = M/6$
$\text{C}_2\text{O}_4^{2-}$	$\text{MnO}_4^-$	$\text{C}_2\text{O}_4^{2-} \longrightarrow 2\text{CO}_2 + 2e^-$ $\text{MnO}_4^- + 8\text{H}^+ + 5e^- \longrightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$	$5\text{C}_2\text{O}_4^{2-} \equiv 2\text{MnO}_4^-$ Eq. wt. $\text{C}_2\text{O}_4^{2-} = M/2$ Eq. wt. $\text{MnO}_4^- = M/5$
$\text{H}_2\text{O}_2$	$\text{MnO}_4^-$	$\text{H}_2\text{O}_2 \longrightarrow 2\text{H}^+ + \text{O}_2 + 2e^-$ $\text{MnO}_4^- + 8\text{H}^+ + 5e^- \longrightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$	$5\text{H}_2\text{O}_2 \equiv 2\text{MnO}_4^-$ Eq. wt. $\text{H}_2\text{O}_2 = M/2$ Eq. wt. $\text{MnO}_4^- = M/5$
$\text{As}_2\text{O}_3$ $\text{AsO}_3^{3-}$	$\text{MnO}_4^-$ $\text{BrO}_3^-$	$\text{As}_2\text{O}_3 + 5\text{H}_2\text{O} \longrightarrow 2\text{AsO}_4^{3-} + 10\text{H}^+ + 4e^-$ $\text{AsO}_3^{3-} + \text{H}_2\text{O} \longrightarrow \text{AsO}_4^{3-} + 2\text{H}^+ + 2e^-$ $\text{BrO}_3^- + 6\text{H}^+ + 6e^- \longrightarrow \text{Br}^- + 3\text{H}_2\text{O}$	Eq. wt. $\text{As}_2\text{O}_3 = M/4$ Eq. wt. $\text{AsO}_3^{3-} = M/2$ Eq. wt. $\text{BrO}_3^- = M/6$



**Ex.** 20 g of a sample of  $\text{Ba(OH)}_2$  is dissolved in 10 mL of 0.5 N HCl solution : The excess of HCl was titrated with 0.2 N NaOH. The volume of NaOH used was 20 cc. Calculate the percentage of  $\text{Ba(OH)}_2$  in the sample.

**Sol.** Milli eq. of HCl initially =  $10 \times 0.5 = 5$

Milli eq. of NaOH consumed = Milli eq. of HCl in excess =  $10 \times 0.2 = 2$

$\therefore$  Milli eq. of HCl consumed = Milli eq. of  $\text{Ba(OH)}_2 = 5 - 2 = 3$

$\therefore$  Eq. of  $\text{Ba(OH)}_2 = 3/1000 = 3 \times 10^{-3}$

Mass of  $\text{Ba(OH)}_2 = 3 \times 10^{-3} \times (171/2) = 0.2565 \text{ g}$

%  $\text{Ba(OH)}_2 = (0.2565 / 20) \times 100 = 1.28 \%$

**Ex.** 3.2 g of pyrolusite was treated with 50 mL of 0.5 M oxalic acid and some sulphuric acid. The oxalic acid left undecomposed was raised to 250 mL in a flask. 25 mL of this solution when treated with 0.02 M  $\text{KMnO}_4$  required 32 mL of the solution : Find the % of  $\text{MnO}_2$  in the sample and also the percentage of available oxygen.

**Sol.** Redox changes are  $\text{C}_2\text{O}_4^{2-} \longrightarrow 2\text{CO}_2$  (n-factor = 2)

$\text{MnO}_4^- \longrightarrow \text{Mn}^{2+}$  (n-factor = 5)

$\text{MnO}_2 \longrightarrow \text{Mn}^{2+}$  (n-factor = 2)

Meq. of  $\text{MnO}_2$  = Meq. of oxalic acid taken - Meq. of oxalic acid left

=  $50 \times 0.5 \times 2 - 32 \times 0.02 \times 5 \times 10$  (in 250 mL) = 18

$\frac{W_{\text{MnO}_2}}{M_{\text{MnO}_2}} \times 2 \times 1000 = 18 \Rightarrow \frac{W_{\text{MnO}_2}}{87} \times 2 \times 1000 = 18,$

$\therefore W_{\text{MnO}_2} = 0.7821 \text{ g}$

$\therefore \% \text{ of MnO}_2 = \frac{0.7821}{3.2} \times 100 = 24.44 \%$

Meq. of  $\text{MnO}_2$  = Meq. of  $\text{O}_2$

$\frac{W_{\text{O}_2}}{16} \times 2 \times 1000 = 18, \therefore W_{\text{O}_2} = 0.144 \text{ g}$


% of available  $\text{O}_2 = \frac{0.144}{3.2} \times 100 = 4.5$

## □ DOUBLE TITRATION



The method involves two indicators (Indicators are substances that change their colour when a reaction is complete) phenolphthalein and methyl orange. This is a titration of specific compounds. Let us consider a solid mixture of NaOH,  $\text{Na}_2\text{CO}_3$  and inert impurities weighing w g. You are asked to find out the % composition of mixture. You are also given a reagent that can react with the sample say, HCl along with its concentration ( $M_1$ )

We first dissolve the mixture in water to make a solution and then we add two indicators in it, namely, phenolphthalein and methyl orange, Now, we titrate this solution with HCl.

NaOH is a strong base while  $\text{Na}_2\text{CO}_3$  is a weak base. So, it is safe to assume that NaOH reacts completely and only then  $\text{Na}_2\text{CO}_3$  reacts.



Once NaOH has reacted, it is the turn of  $\text{Na}_2\text{CO}_3$  to react. It reacts with HCl in two steps :



As can be seen, when we go on adding more and more of HCl, the pH of the solution keeps on falling. When  $\text{Na}_2\text{CO}_3$  is converted to  $\text{NaHCO}_3$ , completely, the solution is weakly basic due to presence of  $\text{NaHCO}_3$  (which is a weaker base as compared to  $\text{Na}_2\text{CO}_3$ ). At this instant phenolphthalein changes colour since it requires this

weakly basic solution to change its colour. Therefore, remember that **phenolphthalein changes colour only when the weakly basic  $\text{NaHCO}_3$  is present**. As we keep adding HCl, the pH again falls and when all the  $\text{NaHCO}_3$  reacts to form NaCl,  $\text{CO}_2$  and  $\text{H}_2\text{O}$ , the solution becomes weakly acidic due to the presence of the weak acid  $\text{H}_2\text{CO}_3$  ( $\text{CO}_2 + \text{H}_2\text{O}$ ). At this instance, methyl orange changes colour since it requires this weakly acidic solution to do so. Therefore, remember **methyl orange changes colour only when  $\text{H}_2\text{CO}_3$  is present**.

Now, let us assume that the volume of HCl used up for the first and the second reaction, i.e.,  $\text{NaOH} + \text{HCl} \rightarrow \text{NaCl} + \text{H}_2\text{O}$  and  $\text{Na}_2\text{CO}_3 + \text{HCl} \rightarrow \text{NaHCO}_3 + \text{NaCl}$  be  $V_1$  (this is the volume of HCl from the beginning of the titration up to the point when phenolphthalein changes colour).

Let, the volume of HCl required for the last reaction, i.e.  $\text{NaHCO}_3 + \text{HCl} \rightarrow \text{NaCl} + \text{CO}_2 + \text{H}_2\text{O}$  be  $V_2$  (this is the volume of HCl from the point where phenolphthalein had changed colour up to the point when methyl orange changes colour). Then,

$$\text{Moles of HCl used for reacting with NaHCO}_3 = \text{Moles of NaHCO}_3 \text{ reacted} = M_1V_2$$

$$\text{Moles of NaHCO}_3 \text{ produced by the Na}_2\text{CO}_3 = M_1V_2$$

$$\text{Moles of Na}_2\text{CO}_3 \text{ that gave } M_1V_2 \text{ moles of NaHCO}_3 = M_1V_2$$

$$\text{Mass of Na}_2\text{CO}_3 = M_1V_2 \quad 106$$

$$\% \text{ Na}_2\text{CO}_3 = \frac{M_1V_2 \times 106}{w} \times 100$$

$$\text{Moles of HCl used for the first two reactions} = M_1V_1$$

$$\text{Moles of Na}_2\text{CO}_3 = M_1V_2$$

$$\text{Moles of HCl used for reacting with Na}_2\text{CO}_3 = M_1V_2$$

$$\text{Moles of HCl used for reacting with only NaOH} = M_1V_1 - M_1V_2$$

$$\therefore \text{Moles of NaOH} = M_1V_1 - M_1V_2$$

$$\text{Mass of NaOH} = (M_1V_1 - M_1V_2) \quad 40$$

$$\% \text{ NaOH} = \frac{(M_1V_1 - M_1V_2) \times 40}{w} \times 100$$

## WORKING RANGE OF FEW INDICATORS

Indicator	pH range	Behaving as
Phenolphthalein	8 – 10	Weak organic acid
Methyl orange	3 – 4.4	Weak organic base

Thus, methyl orange with lower pH range can indicate complete neutralization of all types of bases. Extent of reaction for different bases with acid (HCl) using these two indicators is summarized below :

	Phenolphthalein	Methyl orange
NaOH	100% reaction is indicated $\text{NaOH} + \text{HCl} \longrightarrow \text{NaCl} + \text{H}_2\text{O}$	100% reaction is indicated $\text{NaOH} + \text{HCl} \longrightarrow \text{NaCl} + \text{H}_2\text{O}$
$\text{Na}_2\text{CO}_3$	50% reaction upto $\text{NaHCO}_3$ stage is indicated $\text{Na}_2\text{CO}_3 + \text{HCl} \longrightarrow \text{NaHCO}_3 + \text{NaCl}$	100% reaction is indicated $\text{Na}_2\text{CO}_3 + 2\text{HCl} \longrightarrow 2\text{NaCl} + \text{H}_2\text{O} + \text{CO}_2$
$\text{NaHCO}_3$	No reaction is indicated	100% reaction is indicated $\text{NaHCO}_3 + \text{HCl} \longrightarrow \text{NaCl} + \text{H}_2\text{O} + \text{CO}_2$

**Ex.** 0.4 g of a mixture of NaOH and Na<sub>2</sub>CO<sub>3</sub> and inert impurities was first titrated with phenolphthalein and N/10 HCl, 17.5 mL of HCl was required at the end point. After this methyl orange was added and 1.5 mL of same HCl was again required for next end point. Find out percentage of NaOH and Na<sub>2</sub>CO<sub>3</sub> in the mixture.

**Sol.** Let W<sub>1</sub> g NaOH and W<sub>2</sub> g Na<sub>2</sub>CO<sub>3</sub> was present in mixture. At phenolphthalein end point,

$$\left( \frac{W_1}{40} + \frac{1}{2} \times \frac{W_2}{53} \right) = \frac{1}{10} \times 17.5 \times 10^{-3} \quad \dots\dots\dots(1)$$

At second end point following reaction takes place

Eq. of NaHCO<sub>3</sub> = Eq. of HCl used (in second titration)

$$= \frac{1}{2} \text{Eq. of Na}_2\text{CO}_3$$

$$\frac{1}{2} \times \frac{W_2}{53} = 1.5 \times \frac{1}{10} \times 10^{-3}$$

$$W_2 = 0.01590 \text{ g}$$

Putting the value of W<sub>2</sub> in Eq. (1) we get



$$W_1 = 0.064 \text{ g}$$

$$\text{Percentage of NaOH} = \frac{0.064}{0.4} \times 100 = 16\%$$

$$\text{Percentage of Na}_2\text{CO}_3 = \frac{0.01590}{0.4} \times 100 = 3.975 \%$$

#### **Iodometric and Iodimetric Titration :**


The reduction of free iodine to iodide ions and oxidation of iodide ions to free iodine occurs in these titrations.






These are divided into two types :

#### **Iodometric Titration :**

In iodometric titrations, an oxidizing agent is allowed to react in neutral medium or in acidic medium with excess of potassium iodide to liberate free iodine.




Free iodine is titrated against a standard reducing agent usually with sodium thiosulphate. Halogen, dichromates, cupric ion, peroxides etc., can be estimated by this method.



#### **Iodimetric Titration**

These are the titrations in which free iodine is used as it is difficult to prepare the solution of iodine (volatile and less soluble in water), it is dissolved in KI solution :



This solution is first standardized before using with the standard solution of substance such as sulphite, thiosulphate, arsenite etc, are estimated.

In iodimetric and iodometric titration, starch solution is used as an indicator. Starch solution gives blue or violet colour with free iodine. At the end point, the blue or violet colour disappears when iodine is completely changed to iodide.

❑ SOME IODOMETRIC TITRATIONS (TITRATING SOLUTIONS IS  $\text{Na}_2\text{S}_2\text{O}_3 \cdot 5\text{H}_2\text{O}$ )

Estimation of	Reaction	Relation between O.A. and R.A.
$\text{I}_2$	$\text{I}_2 + 2\text{Na}_2\text{S}_2\text{O}_3 \longrightarrow 2\text{NaI} + \text{Na}_2\text{S}_4\text{O}_6$ <p style="text-align: center;">or</p> $\text{I}_2 + 2\text{S}_2\text{O}_3^{2-} \longrightarrow 2\text{I}^- + \text{S}_4\text{O}_6^{2-}$	$\text{I}_2 = 2\text{I} \equiv 2\text{Na}_2\text{S}_2\text{O}_3$ <p style="text-align: center;">Eq. wt. of <math>\text{Na}_2\text{S}_2\text{O}_3 = \text{M}/1</math></p>
$\text{CuSO}_4$	$2\text{CuSO}_4 + 4\text{KI} \longrightarrow 2\text{Cu}_2\text{I}_2 + 2\text{K}_2\text{SO}_4 + \text{I}_2$ $\text{Cu}^{2+} + 4\text{I}^- \longrightarrow \text{Cu}_2\text{I}_2 + \text{I}_2$ <p style="text-align: center;">(White ppt.)</p> $\text{CaOCl}_2 + \text{H}_2\text{O} \longrightarrow \text{Ca(OH)}_2 + \text{Cl}_2$	$2\text{CuSO}_4 \equiv \text{I}_2 \equiv 2\text{I} \equiv 2\text{Na}_2\text{S}_2\text{O}_3$ <p style="text-align: center;">Eq. wt. of <math>\text{CuSO}_4 = \text{M}/1</math></p>
$\text{CaOCl}_2$	$\text{Cl}_2 + 2\text{KI} \longrightarrow 2\text{KCl} + \text{I}_2$ $\text{Cl}_2 + 2\text{I}^- \longrightarrow 2\text{Cl}^- + \text{I}_2$ $\text{MnO}_2 + 4\text{HCl (conc)} \xrightarrow{\Delta} \text{MnCl}_2 + \text{Cl}_2 + 2\text{H}_2\text{O}$ $\text{Cl}_2 + 2\text{KI} \longrightarrow 2\text{KCl} + \text{I}_2$	$\text{CaOCl}_2 \equiv \text{Cl}_2 \equiv \text{I}_2 \equiv 2\text{I} \equiv 2\text{Na}_2\text{S}_2\text{O}_3$ <p style="text-align: center;">Eq. wt. of <math>\text{CaOCl}_2 = \text{M}/2</math></p>
$\text{MnO}_2$	<p style="text-align: center;">or</p> $\text{MnO}_2 + 4\text{H}^+ + 2\text{Cl}^- \longrightarrow \text{Mn}^{2+} + 2\text{H}_2\text{O} + \text{Cl}_2$ $\text{Cl}_2 + 2\text{I}^- \longrightarrow \text{I}_2 + 2\text{Cl}^-$	$\text{MnO}_2 \equiv \text{Cl}_2 \equiv \text{I}_2 \equiv 2\text{I} \equiv 2\text{Na}_2\text{S}_2\text{O}_3$ <p style="text-align: center;">Eq. wt. of <math>\text{MnO}_2 = \text{M}/2</math></p>
$\text{IO}_3^-$	$\text{IO}_3^- + 5\text{I}^- + 6\text{H}^+ \longrightarrow 3\text{I}_2 + 3\text{H}_2\text{O}$	$\text{IO}_3^- \equiv 3\text{I}_2 \equiv 6\text{I} \equiv 6\text{Na}_2\text{S}_2\text{O}_3$ <p style="text-align: center;">Eq. wt. of <math>\text{IO}_3^- = \text{M}/6</math></p>
$\text{H}_2\text{O}_2$	$\text{H}_2\text{O}_2 + 2\text{I}^- + 2\text{H}^+ \longrightarrow \text{I}_2 + 2\text{H}_2\text{O}$	$\text{H}_2\text{O}_2 \equiv \text{I}_2 \equiv 2\text{I} \equiv 2\text{Na}_2\text{S}_2\text{O}_3$ <p style="text-align: center;">Eq. wt. of <math>\text{H}_2\text{O}_2 = \text{M}/2</math></p>
$\text{Cl}_2$	$\text{Cl}_2 + 2\text{I}^- \longrightarrow 2\text{Cl}^- + \text{I}_2$	$\text{Cl}_2 \equiv \text{I}_2 \equiv 2\text{I} \equiv 2\text{Na}_2\text{S}_2\text{O}_3$ <p style="text-align: center;">Eq. wt. of <math>\text{Cl}_2 = \text{M}/2</math></p>
$\text{O}_3$	$\text{O}_3 + 6\text{I}^- + 6\text{H}^+ \longrightarrow 3\text{I}_2 + 3\text{H}_2\text{O}$	$\text{O}_3 \equiv 3\text{I}_2 \equiv 6\text{I} \equiv 6\text{Na}_2\text{S}_2\text{O}_3$ <p style="text-align: center;">Eq. wt. of <math>\text{O}_3 = \text{M}/6</math></p>
$\text{Cr}_2\text{O}_7^{2-}$	$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{I}^- \longrightarrow 3\text{I}_2 + 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$	$\text{Cr}_2\text{O}_7^{2-} \equiv 3\text{I}_2 \equiv 6\text{I}$ <p style="text-align: center;">Eq. wt. of <math>\text{Cr}_2\text{O}_7^{2-}</math></p>
$\text{MnO}_4^-$	$2\text{MnO}_4^- + 10\text{I}^- + 16\text{H}^+ \longrightarrow 2\text{MnO}_4^- + 5\text{I}_2 + 8\text{H}_2\text{O}$	$2\text{MnO}_4^- \equiv 5\text{I}_2 \equiv 10\text{I}$ <p style="text-align: center;">Eq. wt. of <math>\text{MnO}_4^- = \text{M}/5</math></p>
$\text{BrO}_3^-$	$\text{BrO}_3^- + 6\text{I}^- + 6\text{H}^+ \longrightarrow \text{Br}^- + 3\text{I}_2 + 3\text{H}_2\text{O}$	$\text{BrO}_3^- \equiv 3\text{I}_2 \equiv \text{I}_2$ <p style="text-align: center;">Eq. wt. of <math>\text{BrO}_3^- = \text{M}/6</math></p>
As(V)	$\text{H}_3\text{AsO}_4 + 2\text{I}^- + 2\text{H}^+ \longrightarrow \text{H}_3\text{AsO}_3 + \text{H}_2\text{O} + \text{I}_2$	$\text{H}_3\text{AsO}_4 \equiv \text{I}_2 \equiv 2\text{I}$ <p style="text-align: center;">Eq. wt. of <math>\text{H}_3\text{AsO}_4 = \text{M}/2</math></p>

❑ SOME IODIMETRIC TITRATION (TITRATING SOLUTIONS IS I<sub>2</sub> IN KI)

Estimation of	Reaction	Relation between O.A. and R.A.
H <sub>2</sub> S (in acidic medium)	$\text{H}_2\text{S} + \text{I}_2 \longrightarrow \text{S} + 2\text{I}^- + 2\text{H}^+$	$\text{H}_2\text{S} \equiv \text{I}_2 \equiv 2\text{I}$ Eq. wt. of H <sub>2</sub> S = M/2
SO <sub>3</sub> <sup>2-</sup> (in acidic medium)	$\text{SO}_3^{2-} + \text{I}_2 + \text{H}_2\text{O} \longrightarrow \text{SO}_3^{2-} + 2\text{I}^- + 2\text{H}^+$	$\text{SO}_3^{2-} \equiv \text{I}_2 \equiv 2\text{I}$ Eq. wt. of SO <sub>3</sub> <sup>2-</sup> = M/2
Sn <sup>2+</sup> (in acidic medium)	$\text{Sn}^{2+} + \text{I}_2 \longrightarrow \text{Sn}^{4+} + 2\text{I}^-$	$\text{Sn}^{2+} \equiv \text{I}_2 \equiv 2\text{I}$ Eq. wt. of Sn <sup>2+</sup> = M/2
As(III) (at pH = 8)	$\text{H}_2\text{AsO}_3^- + \text{I}_2 + \text{H}_2\text{O} \longrightarrow \text{HAsO}_4^{2-} + 2\text{I}^- + 2\text{H}^+$	$\text{H}_2\text{AsO}_3^- \equiv \text{I}_2 \equiv 2\text{I}$ Eq. wt. of H <sub>2</sub> AsO <sub>3</sub> <sup>-</sup> = M/2
N <sub>2</sub> H <sub>4</sub>	$\text{N}_2\text{H}_4 + 2\text{I}_2 \longrightarrow \text{N}_2 + 4\text{H}^+ + 4\text{I}^-$	$\text{N}_2\text{H}_4 \equiv 2\text{I}_2 \equiv 4\text{I}$ Eq. wt. of N <sub>2</sub> H <sub>4</sub> = M/4

❑ CONCEPTS AND FORMULA AT A GLANCE

- Number of moles of molecules =  $\frac{\text{wt. in g}}{\text{Mol. wt.}}$

Number of moles of atoms =  $\frac{\text{wt. in g}}{\text{Atomic mass}}$

Number of moles of gases =  $\frac{\text{Volume at STP}}{\text{Standard molar volume}}$

Number of moles of particles, e.g. atoms, molecular ions etc. =  $\frac{\text{Number of particles}}{\text{Avogadro No.}}$

Moles of solute in solution = M V(L)
- Equivalent wt. of element =  $\frac{\text{Atomic wt.}}{\text{Valence}}$

Equivalent wt. of compound =  $\frac{\text{Mol. wt.}}{\text{Total charge on cation or anion}}$

Equivalent wt. of acid =  $\frac{\text{Mol. wt.}}{\text{Basicity}}$

Equivalent wt. of base =  $\frac{\text{Mol. wt.}}{\text{Acidity}}$

Equivalent wt. of an ion =  $\frac{\text{Formula wt.}}{\text{Charge on ion}}$

Equivalent wt. of acid salt =  $\frac{\text{Molecular wt.}}{\text{Replaceable H atom in acid salt}}$

Equivalent wt. of oxidizing or reducing agent =  $\frac{\text{Mol. wt.}}{\text{Change in oxidation number per mole}}$

No. of equivalent = N V(L) =  $\frac{\text{wt. in g}}{\text{Eq. wt.}}$

$$3. \quad \text{Molarity (M)} = \frac{W_s \times 1000}{M_s \times V} = \frac{x \times d \times 10}{M_s}$$

where

$W_s$  = wt. of solute in g

$M_s$  = Mol. wt. of solute

$x$  = % by mass of solute

$d$  = density of solution in g/ml

$V$  = volume of solution in ml

$$4. \quad \text{Normality (N)} = \frac{W_s \times 1000}{E_s \times V} = \frac{x \times d \times 1000}{E_s}$$

$W_s$  = wt. of solute in g.

$E_s$  = eqv. wt. of solute

$V$  = volume of solution in ml

$x$  = % by mass of solute

$d$  = density of solution in g/ml

$$5. \quad \text{Moles} = M \quad V(L) = \frac{\text{wt. of solute}}{\text{Mol. wt.}}$$

$$\text{Millimoles} = M \quad V(\text{ml}) = \frac{\text{wt. of solute}}{\text{Mol. wt.}} \times 1000$$

$$\text{Equivalents of solute} = N \quad V(L)$$

$$\text{Meq. of solute} = \frac{\text{wt.}}{\text{Eq. wt.}} \times 1000$$

6. Molarity equation :

If a solution having molarity  $M_1$  and volume  $V_1$  is diluted to volume  $V_2$  so that new molarity is  $M_2$  then total number of moles remains the same.

$$M_1 V_1 = M_2 V_2$$

For a balanced equation involving  $n_1$  moles of reactant 1 and  $n_2$  moles of reactant 2.

$$\frac{M_1 V_1}{n_1} = \frac{M_2 V_2}{n_2}$$

Normality equation : According to the law of equivalence, the substances combine together in the ratio

$$\text{of their equivalent masses} \quad \frac{\text{wt. of A}}{\text{wt. of B}} = \frac{\text{Eq. wt. of A}}{\text{Eq. wt. of B}}$$

$$\Rightarrow \frac{\text{wt. of A}}{\text{Eq. wt. of A}} = \frac{\text{wt. of B}}{\text{Eq. wt. of B}}$$

Number of gram equivalents of A = Number of gram equivalents of B

$$\text{Number of gram equivalents of A} = \frac{N_A \times V_A}{1000}$$

$$\text{Number of gram equivalents of B} = \frac{N_B V_B}{1000}$$

$$\Rightarrow \frac{N_A V_A}{1000} = \frac{N_B V_B}{1000}$$

$$\Rightarrow N_A V_A = N_B V_B$$

The above equation is called normality equation.

7. Normality (N) = Molarity (M)  $\times$  n  
 (where n = n-factor)  
 For acid-base (neutralization reaction or redox reaction)  
 $N_1V_1 = N_2V_2$  always true  
 But  $M_1V_1 = M_2V_2$  (may or may not be true)  
 But  $M_1n_1V_1 = M_2n_2V_2$  (always true where n-terms represent n-factor)
8. Molality (m) =  $\frac{\text{Moles of solute}}{\text{wt. of solvent (in kg)}}$
9. Strength of solution (S) = N  $\times$  Eq.wt.  
 = M  $\times$  mol. wt.

### SOLVED PROBLEMS (SUBJECTIVE)

**Ex 1.** What volume of 0.40 M  $\text{H}_2\text{SO}_4$  is required to produce 34.0 g of  $\text{H}_2\text{S}$  by the reaction,



**Sol.** 1 mole of  $\text{H}_2\text{S} \equiv 5$  moles of  $\text{H}_2\text{SO}_4$

34 g of  $\text{H}_2\text{S} \equiv 5$  moles of  $\text{H}_2\text{SO}_4$

$$0.40 \quad V_{\text{H}_2\text{SO}_4} = 5$$


$$\therefore V_{\text{H}_2\text{SO}_4} = \frac{5}{0.40} = 12.5 \text{ litre}$$

**Ex 2.** To 50 L of 0.2 N NaOH, 2.5 L of 2N HCl and 15 L of 0.1 N  $\text{FeCl}_3$  solutions are added. What weight of  $\text{Fe}_2\text{O}_3$  can be obtained from the precipitate? Also report the normality of NaOH left in resultant solution :

**Sol.** Eq. of NaOH = 50  $\times$  0.2 = 10

Eq. of HCl = 2.5  $\times$  2 = 5

Eq. of NaOH left after reaction with HCl = 10 - 5 = 5



$\text{FeCl}_3$  reacts with NaOH to give  $\text{Fe}(\text{OH})_3$  which on ignition gives  $\text{Fe}_2\text{O}_3$

$\therefore$  Eq. of NaOH used for  $\text{FeCl}_3$  = Eq. of  $\text{Fe}(\text{OH})_3$

$$= \text{Eq. of } \text{Fe}_2\text{O}_3$$

$$= 15 \times 0.1 = 3.5$$

$\therefore$  Eq. of NaOH left finally = 5 - 3.5 = 1.5



$$\text{Normality of NaOH in the resultant solution} = \frac{3.5}{70} = 0.05$$

$$\frac{W_{\text{Fe}_2\text{O}_3}}{M_{\text{Fe}_2\text{O}_3}} \times 6 = 1.5 \quad (\text{n-factor for } \text{Fe}_2\text{O}_3 = 6)$$

$$W_{\text{Fe}_2\text{O}_3} = \frac{1.5 \times 160}{6} = 40\text{g}$$

**Ex 3.** One litre of a mixture of  $O_2$  and  $O_3$  at NTP was allowed to react with an excess of acidified solution of KI. The iodine liberated required 30 mL of M/10  $Na_2S_2O_3$  solution for titration. What is the weight percent of ozone in the mixture ?

Ultraviolet radiation of wavelength 300 nm can decompose ozone. Assuming that one photon can decompose one ozone molecule, how many photons would have been required for the complete decomposition of ozone in the original mixture?



$$\begin{aligned} \text{Millimoles of } O_3 &= \text{Millimoles of } I_2 = \frac{1}{2} \quad \text{Millimoles of } Na_2S_2O_3 \\ &= \frac{1}{2} \quad 30 \quad 1/10 = 1.5 \text{ millimoles} \\ &= 0.0015 \text{ moles} \end{aligned}$$

Total moles of  $O_3$  and  $O_2$  in the mixture

$$PV = nRT$$

$$1 \times 1 = n \times 0.0821 \times 273$$

$$\therefore n = 0.044$$

$$\text{Moles of } O_2 = 0.044 - 0.0015 = 0.0425$$

$$W_{O_2} = 0.0425 \times 32 \text{ g} = 1.36 \text{ g}$$

$$W_{O_3} = 0.0015 \times 48 \text{ g} = 0.072 \text{ g}$$

$$\therefore \% \text{ of } O_3 = \frac{0.072}{1.432} \times 100 = 5.028 \%$$

$$\text{No. of photons required for decomposition of Ozone molecules} = 0.0015 \times 6.023 \times 10^{23} = 0.90 \times 10^{21}$$

**Ex 4.** A 20 g sample of only CuS and  $Cu_2S$  was treated with 100 mL of 1.25 M  $K_2Cr_2O_7$ . The products obtained were  $Cr^{3+}$ ,  $Cu^{2+}$  and  $SO_2$ . The excess oxidant was reacted with 50 mL of  $Fe^{2+}$  Solution : 25 mL of the same  $Fe^{2+}$  solution required 0.875 M  $KMnO_4$  under acidic condition, the volume of  $KMnO_4$  used was 20 mL. Find the % of CuS and  $Cu_2S$  in the sample.

**Sol.** Equivalents of dichromate initially =  $\frac{1.25 \times 6 \times 100}{1000} = 0.75$

$$\text{Equivalents of } Fe^{2+} \text{ in 25 mL} = \frac{0.875 \times 5 \times 5 \times 20}{1000} = 0.0875$$

$$\text{Equivalents of } Fe^{2+} \text{ in 50 mL} = 0.0875 \times 2 = 0.175$$

$$\text{Equivalents of excess dichromate} = 0.175$$

$$\therefore \text{Equivalents of dichromate consumed by (CuS and } Cu_2S) = 0.75 - 0.175 = 0.575$$

If x g is the mass of CuS, the mass of  $Cu_2S$  is (20 - x) g

$$\frac{x}{95.5} \times 6 + \frac{(20-x)}{159} \times 8 = 0.575$$

$$\therefore x = 5.74 \text{ g}$$

$$\% \text{ CuS} = \frac{5.74}{20} \times 100 = 28.7 \%$$



$$\% \text{ } Cu_2S = 71.3 \%$$



**Ex 5.** 2.249 g of a sample of pure  $\text{BaCO}_3$  and impure  $\text{CaCO}_3$  containing some  $\text{CaO}$  was treated with dil.  $\text{HCl}$  and it evolved 168 mL of  $\text{CO}_2$  at N.T.P. From this solution,  $\text{BaCrO}_4$  was precipitated filtered and washed. The precipitate was dissolved in dil.  $\text{H}_2\text{SO}_4$  and diluted to 100 mL. 10 mL of this solution when treated with  $\text{KI}$  solution liberated iodine which required exactly 20 mL of 0.05 N  $\text{Na}_2\text{S}_2\text{O}_3$ . Calculate the percentage of  $\text{CaO}$  in the sample.

**Sol.** Let, weight of  $\text{BaCO}_3$ ,  $\text{CaCO}_3$  and  $\text{CaO}$  are x,y and z respectively.

$$\therefore x + y + z = 2.249$$



$$\text{Meq. of BaCO}_3 = \text{Meq. of BaCrO}_4 = \text{Meq. of I}_2$$

$$\therefore \frac{x}{197/3} \cdot 1000 = 20 \cdot 0.05 \cdot \frac{100}{10}$$

$$\therefore x = 0.657 \text{ g} \quad \dots(1)$$

The equivalent weight of  $\text{BaCrO}_4$  is  $M/3$ , therefore for  $\text{BaCO}_3$ , it should be  $M/3$  also because mole ratio of  $\text{BaCO}_3$  and  $\text{BaCrO}_4$  is 1 : 1.

Applying POAC for C atom,

$$\text{Moles of C in BaCO}_3 + \text{Moles of C in CaCO}_3 = \text{Moles of C in CO}_2$$

$$\therefore \frac{x}{197} + \frac{y}{100} = \frac{168}{22400}$$

$$\Rightarrow 200x + 294y = 295.5 \quad \dots(2)$$

From equation (1) and (2)

$$y = 0.416 \text{ g}$$

$$\therefore 0.657 + 0.416 + z = 2.249$$

$$z = 1.176$$

$$\% \text{ of CaO} = \frac{1.176}{2.249} = 52.29 \%$$

**Ex 6.** For estimating ozone in the air, a certain volume of air is passed through an alkaline  $\text{KI}$  solution when  $\text{O}_2$  is evolved and iodide is oxidized to iodine. When such a solution is acidified, free iodine is evolved which can be titrated with standard  $\text{Na}_2\text{S}_2\text{O}_3$  Solution : In an experiment, 10 L of air at 1 atm and 27 C were passed through an alkaline  $\text{KI}$  solution, and at the end, the iodine was entrapped in a solution which on titration as above required 1.5 mL of 0.02 N  $\text{Na}_2\text{S}_2\text{O}_3$  solution. Calculate volume percentage of ozone in the sample.

**Sol.** The chemical reaction is,  $\text{H}_2\text{O} + \text{KI} + \text{O}_3 \longrightarrow \text{I}_2 + \text{O}_2 + \text{KOH}$

Milliequivalents of iodine = Milliequivalents of  $\text{KI}$  = Milliequivalents of  $\text{O}_3$  reacted

$$\text{Milliequivalents of Na}_2\text{S}_2\text{O}_3 = 1.5 \cdot 0.02 = 3 \cdot 10^{-2}$$

$$\text{Millimoles of iodine} = \frac{3 \times 10^{-2}}{2} = 1.5 \cdot 10^{-2} \quad [\because \text{n-factor for iodine} = 2]$$


$$\text{Millimoles of ozone} = 1.5 \cdot 10^{-2}$$

$$\text{Volume of ozone} = \frac{nRT}{P} = \frac{1.5 \times 10^{-5} \times 0.0821 \times 300}{1} = 36.945 \cdot 10^{-5} \text{ litre}$$

$$\text{Volume percent of ozone} = \frac{36.945 \times 10^{-5}}{10} \cdot 100 = 3.6945 \cdot 10^{-3}$$

**Ex 7.** A 2.18 g sample contains a mixture of XO and  $X_2O_3$ . It reacts with 0.015 moles of  $K_2Cr_2O_7$  to oxidize the sample completely to form  $XO_4^-$  and  $Cr^{3+}$ . If 0.0187 mole of  $XO_4^-$  is formed, what is the atomic mass of X ?

**Sol.**  $XO + K_2Cr_2O_7 \longrightarrow Cr^{3+} + XO_4^-$



Let, wt. of XO in the mixture be x g

Equivalent of  $K_2Cr_2O_7$  consumed by the mixture = 0.015 6

$$\text{Equivalents of } XO = \frac{x}{x+16} \times 5$$

$$\text{Equivalents of } X_2O_3 = \frac{2.18-x}{2x+48} \times 8$$

$$\therefore \frac{x}{x+16} \times 5 + \frac{2.18-x}{2x+48} \times 8 = 0.015 \quad 6$$

Since 1 mole of XO gives 1 mole  $XO_4^-$  and 1 mole of  $X_2O_3$  gives 2 moles of  $XO_4^-$ ,

$$\therefore \frac{x}{x+16} + \frac{2x(2.18-x)}{2x+48} = 0.0187$$

On solving, x = 99

**Ex 8.** An aqueous solution containing 0.10 g  $KIO_3$  (formula wt. = 214.0) was treated with an excess of KI Solution : The solution was acidified with HCl. The liberated  $I_2$  consumed 55 mL of thiosulphate solution to decolourize the blue starch-iodine complex. Calculate molarity of the sodium thiosulphate solution :

**Sol.** Moles of  $KIO_3 = \frac{0.1}{214} = 0.00047$

$$\therefore \text{Moles of } I_2 \text{ liberated from } KIO_3 = \frac{0.00047}{2} = 0.000235$$


Moles of KI reacting = 0.00047 5 = 0.00235 ( $\because$  n-factor for  $KIO_3$  and KI are 5 and 1 respectively)

$$\text{Moles of } I_2 \text{ produced from KI} = \frac{0.00235}{2} = 0.001175$$

Total moles of  $I_2$  produced and reacted = 0.000235 + 0.001175 = 0.00141 equivalents of  $I_2$  reacted = 0.00141 2 = 0.00282 = equivalents of thiosulphate Solution :

$$\text{Molarity} = \frac{0.00282 \times 1000}{55} = 0.0512 \text{ M (For thiosulphate n-factor = 1)}$$

**Ex 9.** Mg can reduce  $NO_3^-$  to  $NH_3$  in basic medium.




A 25.0 mL sample of  $NO_3^-$  solution was treated with Mg. The  $NH_3(g)$  was passed into 100 mL of 0.15 N HCl. The excess of HCl required 32.10 mL of 0.10 N NaOH for neutralization. What was the molarity of  $NO_3^-$  ions in the original sample ?

**Sol.** Meq. of  $NH_3$  formed = Meq. of HCl used  
 $= 100 \quad 0.15 - 32.10 \quad 0.10$   
 $= 11.79$

Here, n-factor of  $NH_3$  is 1 (acid - base reaction)

For redox change,



$$\therefore \text{Meq. of } NH_3 \text{ for n-factor } 8 = 8 \quad 11.79$$

$$\therefore \text{Normality of } NO_3^- = \frac{94.32}{25} = 3.77$$


$$\text{Molarity of } NO_3^- = \frac{3.77}{8} = 0.47125$$

**Ex10.** 30 mL of a solution containing 9.15 g/litre of an oxalate  $K_xH_y(C_2O_4)_z \cdot nH_2O$  are required for titrating 27 mL of 0.12 N NaOH and 36 mL of 0.12 N  $KMnO_4$  separately. Calculate x,y,z and n. Assume all H atoms (except  $H_2O$ ) are replaceable and x,y,z are in the simple ratio of g atoms.

**Sol.** Let molecular weight of oxalate salt is M

(i) n - factor in acid - base reaction = 2

(ii) n - factor in redox titration =  $2z$



$\therefore$  Meq. of acid in 30 mL = Meq. of NaOH used

$$30 \times \frac{9.15}{M} \times y = 27 \times 0.12 \quad \dots(1)$$

$$\text{Also, } 30 \times \frac{9.15}{M} \times (2z) = 36 \times 0.12 \quad \dots(2)$$

$$\text{From equations (1) and (2) } \frac{y}{2z} = \frac{27}{36} \Rightarrow \frac{y}{z} = \frac{3}{2} \quad \dots(3)$$

Also, total cationic charge = total anionic charge

$$\therefore x + y = 2z \quad \dots(4)$$

By equations (3) and (4)

$$x : y : z :: 1 : 3 : 2$$

These are in simplest ratio and molecular formula is  $KH_3(C_2O_4)_2 \cdot nH_2O$

Molecular weight of salt =  $39 + 3 + 176 + 18n = 218 + 18n$

$$\text{From equation (1), } M = \frac{30 \times 9.15 \times 3}{27 \times 0.12} = 254.16$$

$$\therefore 218 + 18n = 254.15$$


$$\therefore n = 2$$

$\therefore$  Oxalate salt is  $KH_3(C_2O_4)_2 \cdot 2H_2O$



**Ex11.** Hydrogen peroxide solution (20 mL) reacts quantitatively with a solution of  $KMnO_4$  (20 mL) acidified with dilute  $H_2SO_4$ . The same volume of the  $KMnO_4$  solution is just decolourized by 10 mL of  $MnSO_4$  in neutral medium simultaneously forming a dark brown precipitate of hydrated  $MnO_2$ . The brown precipitate is dissolved in 20 mL of 0.2 M sodium oxalate under boiling condition in the presence of dilute  $H_2SO_4$ . Write the balanced equations involved in the reactions and calculate the volume strength of  $H_2O_2$ .

**Sol.** Meq. of  $MnO_2$  = Meq. of  $Na_2C_2O_4$

$$= 20 \times 0.2 \times 2 = 8$$



$$\text{Millimoles of } MnO_2 = \frac{8}{2} = 4$$




(Mole ratio is reciprocal of n-factor ratio)

$$\frac{\text{Millimoles of MnO}_4^-}{\text{Millimoles of MnO}_2} = \frac{2}{5}$$

$$\text{Millimoles MnO}_4^- = \frac{2}{5} \quad \text{Millimoles of MnO}_2$$

$$= \frac{2}{5} \quad 4 = \frac{8}{5}$$



$$\frac{\text{Millimoles of H}_2\text{O}_2}{\text{Millimoles of MnO}_4^-} = \frac{5}{2}$$

$$\therefore \text{Millimoles of H}_2\text{O}_2 = \frac{5}{2} \times \frac{8}{5} = 4$$


$$N_{\text{H}_2\text{O}_2} \quad 20 = 4 \quad 2 \quad (\text{n-factor for H}_2\text{O}_2 = 2)$$

$$\therefore N_{\text{H}_2\text{O}_2} = 0.4$$

$$\begin{aligned} \text{Volume strength of H}_2\text{O}_2 &= 5.6 \quad N_{\text{H}_2\text{O}_2} \\ &= 5.6 \quad 0.4 = 2.24 \end{aligned}$$

**Ex12.** 1.0 g of moist sample of mixture of potassium chlorate ( $\text{KClO}_3$ ) and potassium chloride ( $\text{KCl}$ ) was dissolved in water and solution made upto 250 mL. This solution was treated with  $\text{SO}_2$  to reduce all  $\text{ClO}_3^-$  to  $\text{Cl}^-$  and excess of  $\text{SO}_2$  was removed by boiling. The total chloride was precipitated as silver chloride. The weight of precipitate was found to be 1.435 g. In another experiment, 25 mL of the original solution was heated with 30 ml 0.2 N  $\text{FeSO}_4$  and unused  $\text{FeSO}_4$  required 37.5 mL of 0.08 N  $\text{KMnO}_4$  solutions.

Calculate the molar ratio of the  $\text{ClO}_3^-$  to the given mixture.



**Sol.**  $\text{ClO}_3^-$  is reduced to  $\text{Cl}^-$  by  $\text{SO}_2$  and  $\text{ClO}_3^-$  is also reduced to  $\text{Cl}^-$  by  $\text{Fe}^{2+}$ , hence  $\text{AgCl}$  is formed due to total  $\text{Cl}^-$

$$\text{Meq. of Fe}^{2+} \text{ initially taken} = 30 \quad 0.2 = 6$$

$$\text{Meq. of Fe}^{2+} \text{ unused} = 37.5 \quad 0.08 = 3$$

$$\therefore \text{Meq. of Fe}^{2+} = 6.0 - 3.0 = 3.0$$

$$\text{Thus, Meq. of ClO}_3^- \text{ in 25 mL} = 3.0$$

$$\text{Moles of ClO}_3^- \text{ in 25 mL} = \frac{3.0}{1000 \times 6} = 0.0005$$



$$\text{O.N.} \quad 5 \quad -1$$

$$\text{Thus, moles of ClO}_3^- \text{ in 25 mL solution} = 0.0005$$

$\text{ClO}_3^-$  is also reduced to  $\text{Cl}^-$  by  $\text{SO}_2$  in first experiment and precipitated as  $\text{AgCl}$ .

$$\text{Thus, Cl}^- \text{ formed from ClO}_3^- = \text{AgCl from ClO}_3^- = 1.435 \text{ g}$$



$$\text{Total AgCl formed both from actual and Cl}^- \text{ from ClO}_3^- = 1.435 \text{ g}$$

$$= \frac{1.435}{143.5} = 0.01 \text{ mol}$$

$$\text{Thus, AgCl formed due to Cl}^- \text{ only} = 0.01 - 0.0005 = 0.0095 \text{ mol}$$

$$\text{Thus, ClO}_3^- \text{ and Cl}^- \text{ are in molar ratio} = \mathbf{1 : 19}$$

**Ex13.** Chile salt peter a source of  $\text{NaNO}_3$  also contains  $\text{NaIO}_3$ . The  $\text{NaIO}_3$  can be used as a source of iodine produced in the following reactions :



One litre of chile salt peter solution containing 6.80 g  $\text{NaIO}_3$ , is treated with stoichiometric quantity of  $\text{NaHSO}_3$ . Now additional amount of same solution is added to the reaction mixture to bring about the second reaction. How many grams of  $\text{NaHSO}_3$  are required in step 1 and what additional volume of chile salt peter must be added in step 2 to bring in complete conversion of  $\text{I}^-$  to  $\text{I}_2$  ?

**Sol.**  $\text{Meq. of NaHSO}_3 = \text{Meq. of NaIO}_3 = N \quad V = \frac{6.8}{198} \quad 6 \quad 1000$



$$\therefore \text{Meq. of NaHSO}_3 = 206.06$$

$$\frac{W_{\text{NaHSO}_3}}{M/2} \quad 1000 = 206.06$$

$$\Rightarrow W_{\text{NaHSO}_3} = \frac{206.06 \times 104}{2 \times 1000} = 10.715 \text{ g}$$

$$\text{Also, Meq. of } \text{I}^- \text{ formed using } n\text{-factor } 6 = 206.06$$

In step 2, n-factor  $\text{I}^-$  is 1 and n-factor for  $\text{IO}_3^-$  is 5



$$\text{Meq. of } \text{I}^- \text{ formed using } n\text{-factor } 1 = \frac{206.06}{6}$$

$$\text{Meq. of NaIO}_3 \text{ used in 2}^{\text{nd}} \text{ step} = \frac{206.06}{6}$$

$$\Rightarrow \frac{6.8}{198} \quad 5 \quad V = \frac{206.06}{6}$$



$$\therefore V_{\text{NaIO}_3} = 199.99 \text{ mL}$$

**Ex14.** 1 g sample of  $\text{AgNO}_3$  is dissolved in 50 mL of water. It is titrated with 50 mL of KI Solution : The AgI precipitated is filtered off. Excess of KI in filtrate is titrated with M/10  $\text{KIO}_3$  in presence of 6 M HCl till all  $\text{I}^-$  converted into  $\text{ICl}$ . It requires 50 ml of M/10  $\text{KIO}_3$  Solution : 20 mL of the same stock solution of KI require 30 mL of M/10  $\text{KIO}_3$  under similar conditions. Calculate % of  $\text{AgNO}_3$  in the sample. The reaction is :



As present in  $\text{AgNO}_3$  is removed as AgI by adding 50 mL of KI solution of which 20 mL requires 30 mL of M/10  $\text{KIO}_3$

$$\text{Meq. of KI in 20 mL} = \text{Meq. of KIO}_3 = 30 \quad \frac{1}{10} \times 4$$



$$\therefore \text{Eq. wt. of KI} = \frac{M}{2}$$

$$\therefore \text{Meq. of KI in 50 mL added to AgNO}_3 = \frac{30 \times 4 \times 50}{10 \times 20} = 30$$

$$\text{Meq. of KI left unused by AgNO}_3 = 50 \quad \frac{1}{10} \times 4 = 20$$

$$\therefore \text{Meq. of KI used for AgNO}_3 = 30 - 20 = 10$$

$$\therefore \text{Meq. of AgNO}_3 = 10$$

$$\frac{W_{\text{AgNO}_3}}{170} \times 2 \times 1000 = 10$$

$$\therefore W_{\text{AgNO}_3} = 0.85 \text{ g} = 85\%$$


$$\frac{\text{Mole of AgNO}_3}{\text{Mole of KI}} = \frac{1}{1}$$

$$\therefore \text{If equivalent weight of KI} = \frac{M}{2}$$

$$\text{Equivalent weight of AgNO}_3 = \frac{M}{2}$$


**Ex15.** A sample of Mg was burnt in air to give a mixture of MgO and Mg<sub>3</sub>N<sub>2</sub>. The ash was dissolved in 60 Meq. of HCl and the resulting solution was back titrated with NaOH. 12 Meq. of NaOH were required to reach the end point. An excess of NaOH was then added and the solution was distilled. The ammonia released was then trapped in 10 Meq. of acid. Back titration of this solution required 6 Meq. of base. Calculate the % of Mg burnt to the nitride.

**Sol.** Let, total moles of Mg used for MgO and Mg<sub>3</sub>N<sub>2</sub> be a and b respectively



$$a \qquad \qquad \qquad 0$$



$$0 \qquad \qquad \qquad a$$



$$b \qquad \qquad \qquad 0$$

$$0 \qquad \qquad \qquad b/3$$

Now a,  $\frac{b}{3}$  moles respectively of MgO and Mg<sub>3</sub>N<sub>2</sub> are present in the mixture.



or solution contains (a) moles of MgCl<sub>2</sub> from MgO and (b) moles of MgCl<sub>2</sub> from Mg<sub>3</sub>N<sub>2</sub> and  $\frac{2b}{3}$  moles of NH<sub>4</sub>Cl.

Now, moles of HCl or Meq. of HCl (monobasic) = 30 - 12 = 48

$$2a + \frac{8b}{3} = 48 \qquad \qquad \qquad \dots\dots\dots(1)$$

Moles of NH<sub>4</sub>Cl formed = Moles of NH<sub>3</sub> liberated = Moles of HCl used for absorbing NH<sub>3</sub>.

$$\frac{2b}{3} = (10 - 6) = 4 \qquad \qquad \qquad \dots\dots\dots(2)$$

$$\text{From (1), } 2a + \frac{8 \times 6}{3} = 48 \text{ or } a = 16$$


$$\therefore \% \text{ of Mg used for Mg}_3\text{N}_2 = \frac{6}{(6+16)} \times 100 = 27.27\%$$

# SOLVED PROBLEMS (OBJECTIVE)

**Ex 1.** A 0.1097 g sample of  $\text{As}_2\text{O}_3$  required 36.10 mL of  $\text{KMnO}_4$  solution for its titration. The molarity of  $\text{KMnO}_4$  solution is.

- (A) 0.02 (B) 0.04 (C) 0.0122 (D) 0.3

**Sol.**



Let, molarity of  $\text{KMnO}_4$  solution be M

$\therefore$  Eq. of  $\text{As}_2\text{O}_3$  = Eq. of  $\text{KMnO}_4$  solution

$$\frac{0.1097}{198} \times 4 = \frac{36.10 \times M \times 5}{1000} \quad \left( \text{Equivalent weight } \text{As}_2\text{O}_3 = \frac{198}{4} \right)$$

Molarity = 0.0122 M

Hence, (C) is the correct answer.

**Ex 2.** In basic medium,  $\text{CrO}_4^{2-}$  oxidize  $\text{S}_2\text{O}_3^{2-}$  to form  $\text{SO}_4^{2-}$  and itself changes to  $\text{Cr(OH)}_4^-$ . How many mL of 0.154 M  $\text{CrO}_4^{2-}$  are required to react with 40 mL of 0.246 M  $\text{S}_2\text{O}_3^{2-}$ ?

- (A) 200 mL (B) 156.4 mL (C) 170.4 mL (D) 190.4 mL

**Sol.**  $40 \times 0.246 \times 8 = V \times 0.154 \times 3$  (Meq. of  $\text{S}_2\text{O}_3^{2-}$  = Meq. of  $\text{CrO}_4^{2-}$ )


$\therefore V = 170.4$  mL

Hence, (C) is the correct answer.

**Ex 3.** 10 mL of 0.4 M  $\text{Al}_2(\text{SO}_4)_3$  is mixed with 20 mL of 0.6 M  $\text{BaCl}_2$ . Concentration of  $\text{Al}^{3+}$  ion in the solution will be.

- (A) 0.266 M (B) 10.3 M (C) 0.1 M (D) 0.25 M


**Sol.**



$$[\text{Al}^{3+}] = \frac{24}{30 \times 3} = 0.266 \text{ M}$$

Hence (A) is the correct answer.

**Ex 4.** The weight of sodium bromate required to prepare 55.5 mL of 0.672 N solution for cell reaction,



- (A) 1.56 g (B) 0.9386 g (C) 1.23 g (D) 1.32 g

**Sol.** Meq. of  $\text{NaBrO}_3 = 55.5 \times 0.672 = 37.296$

Let weight of  $\text{NaBrO}_3 = W$


$$\therefore \frac{W}{M_{\text{NaBrO}_3}} \times 6 \times 1000 = 37.296 \quad (\text{equivalent weight} = M/6) \text{ of n-factor} = 6$$

$$\therefore \frac{W}{151} \times 6 \times 1000 = 37.296$$

$$\therefore W = 0.9386 \text{ g}$$

Hence, (B) is the correct answer.

**Ex 5.**  $\text{NaIO}_3$  reacts with  $\text{NaHSO}_3$  according to equation



The weight of  $\text{NaHSO}_3$  required to react with 100 mL of solution containing 0.68 g of  $\text{NaIO}_3$  is

- (A) 5.2 g (B) 0.2143 g (C) 2.3 g (D) none of the above

**Sol.**  $\text{Meq. of NaHSO}_3 = \text{Meq. of NaIO}_3 = N \quad V = \frac{0.68}{198} \quad 6 \quad 1000 \quad (\text{I}^{5+} + 6\text{e}^- \rightarrow \text{I}^-)$

$$\therefore \frac{W_{\text{NaHSO}_3}}{M_{\text{NaHSO}_3}} \quad 2 \quad 1000 = \frac{0.68}{198} \quad 6 \quad 100$$


$$W_{\text{NaHSO}_3} = \frac{0.68 \times 6 \times 100 \times 104}{198 \times 1000} = 0.2143$$

Hence (B) is the correct answer.

**Ex 6.** If 0.5 moles of  $\text{BaCl}_2$  is mixed with 0.1 moles of  $\text{Na}_3\text{PO}_4$ , the maximum amount of  $\text{Ba}_3(\text{PO}_4)_2$  that can be formed is

- (A) 0.7 mol (B) 0.5 mol (C) 0.2 mol (D) 0.05 mol

**Sol.** Let us first solve this problem by writing the complete balanced reaction.



We can see that the moles of  $\text{BaCl}_2$  used are  $\frac{3}{2}$  times the moles of  $\text{Na}_3\text{PO}_4$ . Therefore, to react with

0.1 mol of  $\text{Na}_3\text{PO}_4$ , the moles of  $\text{BaCl}_2$  required would be  $0.1 \times \frac{3}{2} = 0.15$ . Since  $\text{BaCl}_2$  is 0.5 mol, we can

conclude that  $\text{Na}_3\text{PO}_4$  is the limiting reagent. Therefore, moles of  $\text{Ba}_3(\text{PO}_4)_2$  formed is  $0.1 \times \frac{1}{2} = 0.05$  mol.

Hence, (D) is the correct answer.

**Ex 7.** 0.52 g of a dibasic acid required 100 mL of 0.2 N NaOH for complete neutralization.

The equivalent weight of acid is

- (A) 26 (B) 52 (C) 104 (D) 156

**Sol.**  $\text{Meq. of Acid} = \text{Meq. of NaOH}$

$$\frac{0.52}{E} \times 1000 = 100 \quad 0.2$$


$$\therefore E = 26$$

Hence (A) is the correct answer.

**Ex 8.** 34 g hydrogen peroxide is present in 1120 mL of Solution : This solution is called

- (A) 10 volume (B) 20 volume (C) 30 volume (D) 32 volume

**Sol.**  $\text{Wt. of H}_2\text{O}_2 \text{ in 1 mL} = \frac{34}{1120} \text{ g}$



34 g of  $\text{H}_2\text{O}_2$  gives 11200 mL of  $\text{O}_2$  at STP

$$\therefore \frac{34}{1120} \text{ g of H}_2\text{O}_2 \equiv \frac{11200}{34} \quad \frac{34}{1120} = 10 \text{ mL of O}_2 \text{ at STP.}$$


Hence, (A) is the correct answer.



**Ex 9.** The number of moles of  $\text{KMnO}_4$  that will be required to react with 2 mol of ferrous oxalate is

- (A)  $\frac{6}{5}$  (B)  $\frac{2}{5}$  (C)  $\frac{4}{5}$  (D) 1

**Sol.**  $\text{Mn}^{7+} + 5 e^- \rightarrow \text{Mn}^{2+}$  ] 3



3 moles of  $\text{KMnO}_4 = 5$  moles of  $\text{FeC}_2\text{O}_4$


$\therefore$  2 mol of ferrous oxalate  $\equiv \frac{6}{5}$  mole of  $\text{KMnO}_4$

Hence, (A) is the correct answer.

**Ex10.** What volume of 0.1 M  $\text{KMnO}_4$  is needed to oxidize 100 mg of  $\text{FeC}_2\text{O}_4$  in acid solution ?

- (A) 4.1 mL (B) 8.2 mL (C) 10.2 mL (D) 4.6 mL

**Sol.** Meq. of  $\text{KMnO}_4 = \text{Meq. of FeC}_2\text{O}_4$



$\therefore V = 4.1 \text{ mL}$

Hence, (A) is the correct answer.


**Ex11.** What volume of 6 M  $\text{HNO}_3$  is needed to oxidize 8 g of  $\text{Fe}^{2+}$  to  $\text{Fe}^{3+}$ ,  $\text{HNO}_3$  gets converted to  $\text{NO}$ ?

- (A) 8 mL (B) 7.936 mL (C) 32 mL (D) 64 mL

**Sol.** Meq. of  $\text{HNO}_3 = \text{Meq. of Fe}^{2+}$

$$\text{or } 6 \times 3 \times V = \frac{8}{56} \times 1000$$

$V = 7.936 \text{ mL}$



Hence, (B) is the correct answer.

**Ex12.** 0.5 g of fuming  $\text{H}_2\text{SO}_4$  (oleum) is diluted with water. This solution is completely neutralized by 26.7 mL of 0.4 N  $\text{KOH}$ . The percentage of free  $\text{SO}_3$  in the sample is

- (A) 30.6% (B) 40.6% (C) 20.6% (D) 50%

**Sol.** Meq. of  $\text{H}_2\text{SO}_4 + \text{Meq. of SO}_3 = \text{Meq. of KOH}$

$$\therefore \frac{(0.5 - x)}{98/2} \times 1000 + \frac{x}{80/2} \times 1000 = 26.7 \times 0.4$$

$\therefore x = 0.103$

$$\therefore \% \text{ of SO}_3 = \frac{0.103}{0.5} \times 100 = 20.6\%$$

Hence, (C) is the correct answer.

**Ex13.** The minimum quantity of  $\text{H}_2\text{S}$  needed to precipitate 63.5 g of  $\text{Cu}^{2+}$  will be nearly.

- (A) 63.5 g                      (B) 31.75 g                      (C) 34 g                      (D) 2.0 g

**Sol.** Meq. of  $\text{H}_2\text{S}$  = Meq. of  $\text{Cu}^{2+}$

$$\therefore \frac{W_{\text{H}_2\text{S}}}{34/2} \times 1000 = \frac{63.5}{63.5/2} \times 1000$$

$$\therefore W_{\text{H}_2\text{S}} = 34 \text{ g}$$

Hence, (C) is the correct answer.

**Ex14.** Which of the following is / are correct?

- (A) g mole wt. = mol. wt. in g = wt. of  $6.02 \times 10^{23}$  molecules  
 (B) mole = N molecule =  $6.02 \times 10^{23}$  molecules  
 (C) mole = g molecules  
 (D) none of the above

**Sol.** (A), (B) and (C)

**Ex15.** 8 g of  $\text{O}_2$  has the same number of molecules as

- (A) 7 g of CO                      (B) 14 g of CO                      (C) 28 g of CO                      (D) 11 g of  $\text{CO}_2$

**Sol.** (A) and (D)

**Ex16.** The eq. wt. of a substance is the weight which either combines or displaces.

- (A) 8 part of O                      (B) 2 part of H                      (C) 35.5 part of Cl                      (D) none of the above

**Sol.** (A) and (C)

**Ex17.** 'A' g of a metal displaces V mL of  $\text{H}_2$  at NTP. Eq. wt of metal E is / are :

$$(A) E = \frac{A}{\text{Wt. of } \text{H}_2 \text{ displaced}} \times \text{Eq. wt. of H} \quad (B) E = \frac{A \times 1.008 \times 22400}{\text{Volume of } \text{H}_2 \text{ displaced} \times 2} \times \text{Eq. wt. of H}$$

$$(C) E = \frac{A \times 1.008}{\text{Volume of } \text{H}_2 \text{ displaced} \times 0.0000897} \quad (D) \text{ none of the above}$$


**Sol.** (A), (B) and (C)

**Ex18.** Which of the following is/are redox reaction (s)?

- (A)  $\text{BaO}_2 + \text{H}_2\text{SO}_4 \rightarrow \text{BaSO}_4 + \text{H}_2\text{O}_2$                       (B)  $2\text{BaO} + \text{O}_2 \rightarrow 2\text{BaO}_2$   
 (C)  $2\text{KClO}_3 \rightarrow 2\text{KCl} + 3\text{O}_2$                       (D)  $\text{SO}_2 + 2\text{H}_2\text{S} \rightarrow 2\text{H}_2\text{O} + 3\text{S}$

**Sol.** (B), (C) and (D)

**Ex19.** In the reaction,



- (A) bromine is oxidized and carbonate is reduced                      (B) bromine is oxidized  
 (C) bromine is reduced                      (D) it is disproportionation reaction or autoredox change

**Sol.** (B), (C) and (D)

**Ex20.** Which of the following statements is/are true if 1 mol of  $\text{H}_3\text{PO}_x$  is completely neutralized by 40 g of NaOH?

- (A) x = 2 and acid is monobasic                      (B) x = 3 and acid is dibasic  
 (C) x = 4 and acid is tribasic                      (D) x = 2 and acid does not form acid salt

**Sol.** (A) and (D).