

RELATIONS

INTRODUCTION :

Let A and B be two sets. Then a relation R from A to B is a subset of $A \times B$.

thus, R is a relation from A to B $\Leftrightarrow R \subseteq A \times B$.

Ex. If $A = \{1, 2, 3\}$ and $B = \{a, b, c\}$, then $R = \{(1, b), (2, c), (1, a), (3, a)\}$ being a subset of $A \times B$, is a relation from A to B. Here $(1, b), (2, c), (1, a)$ and $(3, a) \in R$, so we write $1Rb, 2Rc, 1Ra$ and $3Ra$. But $(2, b) \notin R$, so we write $2 \not R b$.

Total Number of Relations : Let A and B be two non-empty finite sets consisting of m and n elements respectively. Then $A \times B$ consists of mn ordered pairs. So, total number of subsets of $A \times B$ is 2^{mn} .

Domain and Range of a relation : Let R be a relation from a set A to a set B. Then the set of all first components or coordinates of the ordered pairs belonging to R is called to domain of R, while the set of all second components or coordinates of the ordered pairs in R is called the range of R.

Thus, $\text{Dom}(R) = \{a : (a, b) \in R\}$
 and, $\text{Range}(R) = \{b : (a, b) \in R\}$

It is evident from the definition that the domain of a relation from A to B is a subset of A and its range is a subset of B.

e.g. Let $A = \{1, 3, 5, 7\}$ and $B = \{2, 4, 6, 8\}$ be two sets and let R be a relation from A to B defined by the phrase " $(x, y) \in R \Leftrightarrow x > y$ ". Under this relation R, we have

$3R2, 5R2, 5R4, 7R2, 7R4$ and $7R6$

i.e. $R = \{(3, 2), (5, 2), (5, 4), (7, 2), (7, 4), (7, 6)\}$

$\therefore \text{Dom}(R) = \{3, 5, 7\}$ and $\text{Range}(R) = \{2, 4, 6\}$

Inverse Relation : Let A, B be two sets and let R be a relation from a set A to a set B. Then the inverse of R, denoted by R^{-1} , is a relation from B to A and is defined by

$$R^{-1} = \{(b, a) : (a, b) \in R\}$$

Clearly, $(a, b) \in R \Leftrightarrow (b, a) \in R^{-1}$

Also, $\text{Dom}(R) = \text{Range}(R^{-1})$ and $\text{Range}(R) = \text{Dom}(R^{-1})$

Illustration 1 :

Let A be the set of first ten natural numbers and let R be a relation on A defined by $(x, y) \in R \Leftrightarrow x + 2y = 10$, i.e. $R = \{(x, y) : x \in A, y \in A \text{ and } x + 2y = 10\}$. Express R and R^{-1} as sets of ordered pairs. Determine also (i) domain of R and R^{-1} (ii) range of R and R^{-1}

Solution :

We have $(x, y) \in R \Leftrightarrow x + 2y = 10 \Leftrightarrow y = \frac{10-x}{2}, x, y \in A$

where $A = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$

Now, $x = 1 \Rightarrow y = \frac{10-1}{2} = \frac{9}{2} \notin A$.

This shows that 1 is not related to any element in A. Similarly we can observe. that 3, 5, 7, 9 and 10 are not related to any element of A under the defined relation

Further we find that :

For $x = 2, y = \frac{10-2}{2} = 4 \in A \quad \therefore (2, 4) \in R$

For $x = 4, y = \frac{10-4}{2} = 3 \in A \quad \therefore (4, 3) \in R$

For $x = 6, y = \frac{10-6}{2} = 2 \in A \quad \therefore (6, 2) \in R$

$$\text{For } x = 8, y = \frac{10-8}{2} = 1 \in A \quad \therefore (8, 1) \in R$$

$$\text{Thus, } R = \{(2, 4), (4, 3), (6, 2), (8, 1)\}$$

$$\Rightarrow R^{-1} = \{(4, 2), (3, 4), (2, 6), (1, 8)\}$$

$$\text{Clearly, } \text{Dom}(R) = \{2, 4, 6, 8\} = \text{Range}(R^{-1})$$

$$\text{and, } \text{Range}(R) = \{4, 3, 2, 1\} = \text{Dom}(R^{-1})$$

Do yourself - 1 :

(i) If $A = \{2, 4, 6, 9\}$ and $B = \{4, 6, 18, 27, 54\}$, $a \in A$, $b \in B$, find the set of ordered pairs such that 'a' is a factor of 'b' and $a < b$.

(ii) Find the domain and range of the relation R given by $R = \{(x, y) : y = x + \frac{6}{x}, \text{ where } x, y \in \mathbb{N} \text{ and } x < 6\}$

TYPES OF RELATIONS :

In this section we intend to define various types of relations on a given set A .

Void Relation : Let A be a set. Then $\phi \subseteq A \times A$ and so it is a relation on A . This relation is called the void or empty relation on A .

Universal Relation : Let A be a set. Then $A \times A \subseteq A \times A$ and so it is a relation on A . This relation is called the universal relation on A .

Identity Relation : Let A be a set. Then the relation $I_A = \{(a, a) : a \in A\}$ on A is called the identity relation on A .

In other words, a relation I_A on A is called the identity relation if every element of A is related to itself only.

e.g. The relation $I_A = \{(1, 1), (2, 2), (3, 3)\}$ is the identity relation on set $A = \{1, 2, 3\}$. But relations $R_1 = \{(1, 1), (2, 2)\}$ and $R_2 = \{(1, 1), (2, 2), (3, 3), (1, 3)\}$ are not identity relations on A , because $(3, 3) \notin R_1$ and in R_2 element 1 is related to elements 1 and 3.

Reflexive Relation : A relation R on a set A is said to be reflexive if every element of A is related to itself.

Thus, R on a set A is not reflexive if there exists an element $a \in A$ such that $(a, a) \notin R$.

e.g. Let $A = \{1, 2, 3\}$ be a set. Then $R = \{(1, 1), (2, 2), (3, 3), (1, 3), (2, 1)\}$ is a reflexive relation on A . But $R_1 = \{(1, 1), (3, 3), (2, 1), (3, 2)\}$ is not a reflexive relation on A , because $2 \in A$ but $(2, 2) \notin R_1$.

Note : Every Identity relation is reflexive but every reflexive relation is not identity.

Symmetric Relation : A relation R on a set A is said to be a symmetric relation iff

$$(a, b) \in R \Rightarrow (b, a) \in R \text{ for all } a, b \in A$$

i.e. $a R b \Rightarrow b R a$ for all $a, b \in A$.

e.g. Let L be the set of all lines in a plane and let R be a relation defined on L by the rule $(x, y) \in R \Leftrightarrow x$ is perpendicular to y . Then R is a symmetric relation on L , because $L_1 \perp L_2 \Rightarrow L_2 \perp L_1$

$$\text{i.e. } (L_1, L_2) \in R \Rightarrow (L_2, L_1) \in R.$$

e.g. Let $A = \{1, 2, 3, 4\}$ and Let R_1 and R_2 be relation on A given by $R_1 = \{(1, 3), (1, 4), (3, 1), (2, 2), (4, 1)\}$ and $R_2 = \{(1, 1), (2, 2), (3, 3), (1, 3)\}$. Clearly, R_1 is a symmetric relation on A . However, R_2 is not so, because $(1, 3) \in R_2$ but $(3, 1) \notin R_2$

Transitive Relation : Let A be any set. A relation R on A is said to be a transitive relation iff

$$(a, b) \in R \text{ and } (b, c) \in R \Rightarrow (a, c) \in R \text{ for all } a, b, c \in A$$

i.e. $a R b$ and $b R c \Rightarrow a R c$ for all $a, b, c \in A$

e.g. On the set N of natural numbers, the relation R defined by $x R y \Rightarrow x$ is less than y is transitive, because for any $x, y, z \in N$

$$x < y \text{ and } y < z \Rightarrow x < z \Rightarrow x R y \text{ and } y R z \Rightarrow x R z$$

e.g. Let L be the set of all straight lines in a plane. Then the relation 'is parallel to' on L is a transitive relation, because from any $\ell_1, \ell_2, \ell_3 \in L$.

$$\ell_1 \parallel \ell_2 \text{ and } \ell_2 \parallel \ell_3 \Rightarrow \ell_1 \parallel \ell_3$$

Antisymmetric Relation : Let A be any set. A relation R on set A is said to be an antisymmetric relation iff

$$(a, b) \in R \text{ and } (b, a) \in R \Rightarrow a = b \text{ for all } a, b \in A$$

e.g. Let R be a relation on the set N of natural numbers defined by

$$x R y \Leftrightarrow 'x \text{ divides } y' \text{ for all } x, y \in N$$

This relation is an antisymmetric relation on N . Since for any two numbers $a, b \in N$

$$a|b \text{ and } b|a \Rightarrow a = b \quad \text{i.e. } a R b \text{ and } b R a \Rightarrow a = b$$

Equivalence Relation : A relation R on a set A is said to be an equivalence relation on A iff

- (i) it is reflexive i.e. $(a, a) \in R$ for all $a \in A$
- (ii) it is symmetric i.e. $(a, b) \in R \Rightarrow (b, a) \in R$ for all $a, b \in A$
- (iii) it is transitive i.e. $(a, b) \in R$ and $(b, c) \in R \Rightarrow (a, c) \in R$ for all $a, b, c \in A$.

e.g. Let R be a relation on the set of all lines in a plane defined by $(\ell_1, \ell_2) \in R \Leftrightarrow$ line ℓ_1 is parallel to line ℓ_2 . R is an equivalence relation.

Note : It is not necessary that every relation which is symmetric and transitive is also reflexive.

PARTIAL ORDER RELATION :

A relation R on set A is said to be a partial order relation on A if

- (i) R is reflexive i.e. $(a, a) \in R, \forall a \in A$
- (ii) R is antisymmetric i.e. $(a, b) \in R \Rightarrow (b, a) \in R$ only Possible When $a = b \quad \forall a, b \in A$
- (iii) R is transitive i.e. $(a, b) \in R$ and $(b, c) \in R \Rightarrow (a, c) \in R \quad \forall a, b, c \in R$

e.g. R be a relation on the set N of natural numbers defined by

$$x R y \Rightarrow 'x \text{ divides } y' \quad \forall x, y \in N \text{ then } R \text{ is a partial order Relation.}$$

Illustration 2 :

Three relation R_1, R_2 and R_3 are defined on set $A = \{a, b, c\}$ as follows :

- (i) $R_1 \{ (a, a), (a, b), (a, c), (b, b), (b, c), (c, a), (c, b), (c, c) \}$
- (ii) $R_2 \{ (a, b), (b, a), (a, c), (c, a) \}$
- (iii) $R_3 \{ (a, b), (b, c), (c, a) \}$

Find whether each of R_1, R_2 and R_3 is reflexive, symmetric and transitive.

Solution :

- (i) Reflexive : Clearly, $(a, a), (b, b), (c, c) \in R_1$. So, R_1 is reflexive on A .

Symmetric : We observe that $(a, b) \in R_1$ but $(b, a) \notin R_1$. So, R_1 is not symmetric on A .

Transitive : We find that $(b, c) \in R_1$ and $(c, a) \in R_1$ but $(b, a) \notin R_1$. So, R_1 is not transitive on A .

- (ii) Reflexive : Since $(a, a), (b, b)$ and (c, c) are not in R_2 . So, it is not a reflexive relation on A .

Symmetric : We find that the ordered pairs obtained by interchanging the components of ordered pairs in R_2 are also in R_2 . So, R_2 is a symmetric relation on A .

Transitive : Clearly $(c, a) \in R_2$ and $(a, b) \in R_2$ but $(c, b) \notin R_2$. So, it is not a transitive relation on R_2 .

- (iii) Reflexive : Since none of $(a, a), (b, b)$ and (c, c) is an element of R_3 . So, R_3 is not reflexive on A .

Symmetric : Clearly, $(b, c) \in R_3$ but $(c, b) \notin R_3$. So, it is not symmetric on A .

Transitive : Clearly, $(b, c) \in R_3$ and $(c, a) \in R_3$ but $(b, a) \notin R_3$. So, R_3 is not transitive on A .

Illustration 3 :

Prove that the relation R on the set Z of all integers defined by

$$(x, y) \in R \Leftrightarrow x - y \text{ is divisible by } n$$

is an equivalence relation on Z .

Solution :

We observe the following properties

Reflexivity : For any $a \in Z$, we have

$$a - a = 0 = 0 \quad n \Rightarrow a - a \text{ is divisible by } n \Rightarrow (a, a) \in R$$

Thus, $(a, a) \in R$ for all $a \in Z$

So, R is reflexive on Z

symmetry : Let $(a, b) \in R$. Then,

$$(a, b) \in R \Rightarrow (a - b) \text{ is divisible by } n$$

$$\Rightarrow a - b = np \text{ for some } p \in Z$$

$$\Rightarrow b - a = n(-p)$$

$$\Rightarrow b - a \text{ is divisible by } n \quad [\because p \in Z \Rightarrow -p \in Z]$$

$$\Rightarrow (b, a) \in R$$

Thus, $(a, b) \in R \Rightarrow (b, a) \in R$ for all $a, b \in Z$

So, R is symmetric on Z .

Transitivity : Let $a, b, c \in Z$ such that $(a, b) \in R$ and $(b, c) \in R$. Then,

$$(a, b) \in R \Rightarrow (a - b) \text{ is divisible by } n$$

$$\Rightarrow a - b = np \text{ for some } p \in Z$$

$$(b, c) \in R \Rightarrow (b - c) \text{ is divisible by } n$$

$$\Rightarrow b - c = nq \text{ for some } q \in Z$$

$$\therefore (a, b) \in R \text{ and } (b, c) \in R$$

$$\Rightarrow a - b = np \text{ and } b - c = nq$$

$$\Rightarrow (a - b) + (b - c) = np + nq$$

$$\Rightarrow a - c = n(p + q)$$

$$\Rightarrow a - c \text{ is divisible by } n$$

$$[\because p, q \in Z \Rightarrow p + q \in Z]$$

$$\Rightarrow (a, c) \in R$$

thus, $(a, b) \in R$ and $(b, c) \in R \Rightarrow (a, c) \in R$ for all $a, b, c \in Z$. so, R is transitive relation in Z .

Illustration 4 :

Show that the relation 'is congruent to' on the set of all triangles in a plane is an equivalence relation.

Solution :

Let S be the set of all triangles in a plane and let R be the relation on S defined by $(\Delta_1, \Delta_2) \in R \Leftrightarrow$ triangle Δ_1 is congruent to triangle Δ_2 . We observe the following properties.

Reflexivity : For each triangle $\Delta \in S$, we have

$$\Delta \cong \Delta \Rightarrow (\Delta, \Delta) \in R \text{ for all } \Delta \in S \Rightarrow R \text{ is reflexive on } S$$

Symmetry : Let $\Delta_1, \Delta_2 \in S$ such that $(\Delta_1, \Delta_2) \in R$. Then, $(\Delta_1, \Delta_2) \in R \Rightarrow \Delta_1 \cong \Delta_2 \Rightarrow \Delta_2 \cong \Delta_1 \Rightarrow (\Delta_2, \Delta_1) \in R$

So, R is symmetric on S

Transitivity : Let $\Delta_1, \Delta_2, \Delta_3 \in S$ such that $(\Delta_1, \Delta_2) \in R$ and $(\Delta_2, \Delta_3) \in R$. Then,

$(\Delta_1, \Delta_2) \in R$ and $(\Delta_2, \Delta_3) \in R \Rightarrow \Delta_1 \cong \Delta_2$ and $\Delta_2 \cong \Delta_3 \Rightarrow \Delta_1 \cong \Delta_3 \Rightarrow (\Delta_1, \Delta_3) \in R$

So, R is transitive on S .

Hence, R being reflexive, symmetric and transitive, is an equivalence relation on S .

Do yourself - 2 :

- (i) Show that the relation R defined on the set N of natural number by $xRy \Leftrightarrow 2x^2 - 3xy + y^2 = 0$, i.e. by $R = \{(x, y); x, y \in N \text{ and } 2x^2 - 3xy + y^2 = 0\}$ is not symmetric but it is reflexive.

ANSWERS FOR DO YOURSELF

1. (i) $\{(2, 4), (2, 6), (2, 18), (2, 54), (6, 18), (6, 54), (9, 18), (9, 27), (9, 54)\}$
 (ii) Domain of $R = \{1, 2, 3\}$, Range of $R = \{7, 5\}$