

Que


Radius of the largest circle which passes through the focus of the parabola $y^2 = 4x$ and contained in it, is

(A) 8

(B) 4

(C) 2

(D) 5



radius of circle is $PC = \sqrt{(2a)^2 + (2at)^2}$

$$(x - 2a - at^2)^2 + y^2 = (2a)^2 + (2at)^2$$

$$(1,0) \left[\begin{aligned} (1 - 2 - t^2)^2 + 0 &= 4 + 4t^2 \\ (1+t^2)^2 &= 4(1+t^2) \Rightarrow 1+t^2 = 4 \\ t^2 &= 3 \end{aligned} \right.$$

$$r = \sqrt{4(1+t^2)} = \sqrt{4 \times 4} = 4$$

(B)

✓

Ques

The locus of the vertices of the parabolas represented by the equation

$$y = \frac{a^3 x^2}{3} + \frac{a^2 x}{2} - 2a$$

(a is a non-zero parameter) is

- A) a circle B) a parabola
C) a straight line
D) $xy = c^2$ form where 'c' is constant

Sol

$$y = \left(\frac{a^3}{3}\right)x^2 + \left(\frac{a^2}{2}\right)x - 2a$$

$$\begin{array}{l|l} A = \frac{a^3}{3} & D = \frac{a^4}{4} + 4 \cdot \frac{a^3}{3} \cdot 2a \\ B = \frac{a^2}{2} & = a^4 \left(\frac{1}{4} + \frac{8}{3} \right) = \frac{35}{12} a^4 \\ C = -2a & \end{array}$$

$$V: \left(-\frac{B}{2A}, -\frac{D}{4A} \right)$$

$\hookrightarrow x \qquad \qquad \hookrightarrow y$

$$x = -\frac{a^2}{2} \cdot \frac{3}{2a^3} = -\frac{3}{4a}$$

$$y = -\frac{35 \cdot a^4}{\frac{12}{4} \cdot 4a^3} \times \frac{3}{4} = -\frac{35 \cdot a}{16}$$

$$xy = \frac{35 \times 3}{64} = \frac{105}{64}$$

①



$$\text{Product} = -\frac{4f}{a} = x_1 \left(-t_1 - \frac{2}{t_1}\right) \cdot \left(\frac{2}{t_1}\right)$$

$$\Rightarrow \frac{2f}{a} = -\left(t_1 + \frac{2}{t_1}\right)$$

$$\frac{4f^2}{a^2} = \underbrace{t_1^2 + \frac{4}{t_1^2}} + 4 = -\left(2 + \frac{4a+2g}{a}\right) + 4$$

$$\begin{array}{l} \downarrow -g \rightarrow x \\ \downarrow -f \rightarrow y \end{array}$$

$$\frac{4y^2}{a^2} = 2 - \frac{4a-2x}{a} = -2 + \frac{2x}{a}$$

$$\boxed{2y^2 = ax - a^2} \quad \text{Q.E.D.}$$

$$2t_1x - (t_1^2 - 1)y - 2at_1 = 0$$

$$T_1(at_1t_2, a(t_1+t_2)) \rightarrow p_1 = \frac{kt_1at_1t_2 - (t_1^2 - 1) \cdot a(t_1+t_2) - 2at_1}{\sqrt{4t_1^2 + (t_1^2 - 1)^2}}$$


$$p_1 = a \frac{t_1^2t_2 - t_1^3 + t_1 - \cancel{t_1^2t_2} + t_2 - 2t_1}{\sqrt{(t_1^2 + 1)^2}}$$

$$= a \frac{t_1^2t_2 - t_1^3 + t_2 - t_1}{t_1^2 + 1} = \frac{a |t_1^2(t_2 - 1) + 1 \cdot (t_2 - 1)|}{(t_1^2 + 1)}$$

$$= a \frac{\cancel{(t_1^2 + 1)} |t_2 - 1|}{\cancel{(t_1^2 + 1)}} = a |t_2 - t_1|$$

///

$p_2 = a |t_2 - t_1| \Rightarrow$ ST is angular bisector. H.P. (i)



$$T_1(a t_1, a(1+t_1^2))$$

$$\begin{aligned} S P &= a(1+t_1^2) \\ S Q &= a(1+t_2^2) \\ \hline S P \cdot S Q &= a^2(1+t_1^2)(1+t_2^2) \end{aligned}$$

$$S T^2 = S P \cdot S Q$$


H.P.

(i)

$$\begin{aligned} S T^2 &= (a t_1 t_2 - a)^2 + (a(t_1 + t_2) - 0)^2 \\ &= a^2((t_1 t_2 - 1)^2 + (t_1 + t_2)^2) \\ &= a^2(t_1^2 t_2^2 + 1 + t_1^2 + t_2^2) \\ &= a^2(1+t_1^2)(1+t_2^2) \end{aligned}$$

(iii)

$$\frac{S T}{S P} = \frac{S Q}{S T} \quad \text{as } \alpha = \alpha$$



$$\Delta S P T \sim \Delta S T Q$$

H.P. (ii)

One

If l, m are variable real numbers such that $3l^2 + 4m^2 - lm - 3m = 0$, then variable line $lx + my = 1$ always touches a fixed parabola,

Vertex of the parabola is

A) $\left(\frac{1}{3}, -\frac{4}{3}\right)$ B) $\left(-\frac{1}{3}, \frac{4}{3}\right)$

C) $\left(\frac{4}{3}, -\frac{1}{3}\right)$ D) $\left(-\frac{4}{3}, \frac{1}{3}\right)$

Focus of the parabola is

A) $\left(\frac{1}{3}, -\frac{1}{3}\right)$ B) $\left(-\frac{1}{3}, \frac{7}{3}\right)$

C) $\left(-\frac{1}{3}, \frac{1}{3}\right)$ D) $\left(-\frac{7}{3}, \frac{1}{3}\right)$

Directrix of the parabola is

A) $3y + 7 = 0$ B) $3y + 4 = 0$

C) $3y - 4 = 0$ D) $3y - 7 = 0$

$$3l^2 + 4m^2 - lm - 3m = 0$$

$$lx + my = 1$$

$$l = \frac{1 - my}{x}$$

① → (B)

② → (C)

③ → (D)

Concept of Envelope

$$y = mx + \frac{a}{m}$$

↓ (h, k)

$$k = mh + \frac{a}{m}$$


$$hm^2 - km + a = 0$$

$$D > 0$$

$$\frac{k^2 - 4ah}{4h^2} > 0$$

$$S_1 > 0$$

$$D = 0$$



$$3l^2 + 4m^2 - 3m - lm = 0$$

$$l = \frac{1-my}{x} \rightarrow 3 \left(\frac{1+m^2y^2-2my}{x^2} \right) + 4m^2 - 3m + m \left(\frac{my-1}{x} \right) = 0.$$

$$3 + 3m^2y^2 - 6my + \frac{4x^2m^2}{x^2} - 3x^2m + \frac{xy m^2}{x} - mx = 0.$$

$$m^2(3y^2 + 4x^2 + xy) - m(6y + 3x^2 + x) + 3 = 0.$$

$$\Rightarrow D=0 \Rightarrow \cancel{36}y^2 + 9x^4 + x^2 + 6x^3 + \cancel{12xy} + 36yx^2 - 12(\cancel{3x^2} + 4x^2 + \cancel{xy}) = 0$$


$$9x^4 + x^2 + 6x^3 + 36x^2y - 48x^2 = 0.$$

$\div x^2$

$$9x^2 + 1 + 6x + 36y - 48 = 0 \quad \div 9$$

$$(3x+1)^2 = -36\left(y - \frac{48}{36}\right)$$

$$\left(x + \frac{1}{3}\right)^2 = -4\left(y - \frac{4}{3}\right)$$




$y = 7/3 \rightarrow$ Directrix

Ques

A series of chords is drawn so that their projections on the straight line which is inclined at an angle α to the axis are of constant length c . Prove that the locus of their middle point is the curve

$$(y^2 - 4ax)(y \cos \alpha + 2a \sin \alpha)^2 + a^2 c^2 = 0$$

Soln.



$$(\cos \alpha) \hat{i} + (\sin \alpha) \hat{j}$$

$$h = \frac{a(t_1^2 + t_2^2)}{2}, \quad k = a(t_1 + t_2)$$

$$k^2 = a^2(t_1^2 + t_2^2 + 2t_1 t_2)$$

$$k^2 = a(2h) + 2a^2 t_1 t_2$$

$$(t_1 - t_2)^2 = t_1^2 + t_2^2 - 2t_1 t_2$$

$$\overline{AB} = a(t_1^2 - t_2^2) \hat{i} + 2a(t_1 - t_2) \hat{j}$$

$$|\overline{AB} \cdot \hat{u}| = c$$

$$|a(t_1^2 - t_2^2) \cos \alpha + 2a(t_1 - t_2) \sin \alpha| = c$$

$$a^2(t_1 - t_2)^2 \left((t_1 + t_2) \cos \alpha + 2 \sin \alpha \right)^2 = c^2$$

$$a^2 (t_1 - t_2)^2 \left((t_1 + t_2) \cos(\alpha) + 2 \sin(\alpha) \right)^2 = c^2$$

$$\cancel{a^2} \left(\cancel{2ah - k^2} \right) \left(\frac{k}{a} \cos(\alpha) + 2 \sin(\alpha) \right)^2 = c^2$$

$$\begin{array}{l} \Downarrow \\ k \rightarrow y \\ h \rightarrow x \end{array}$$


$$(4ax - y^2) (y \cos(\alpha) + 2a \sin(\alpha))^2 = a^2 c^2$$

$$(y^2 - 4ax) (y \cos(\alpha) + 2a \sin(\alpha))^2 + a^2 c^2 = 0.$$

$$\begin{aligned} (t_1 - t_2)^2 &= (t_1^2 + t_2^2) - 2t_1 t_2 \\ &= \frac{2h}{a} - \cancel{2} \left(\frac{k^2 - 2ah}{\cancel{2}a^2} \right) \\ &= \frac{2ah - k^2 + 2ah}{a^2}. \end{aligned}$$

Qwe

A line AB makes intercepts of length a and b on the coordinate axes. Find the equation of the parabola passing through A, B and the origin if AB is the shortest focal chord of the parabola. Given A, B lie on X-axis & Y-axis resp.



Shortest focal chord
 $\Rightarrow AB$ is L.R.

$$A = \sqrt{a^2 + b^2}$$

$$2A = \frac{\sqrt{a^2 + b^2}}{2}$$

Distance b/w L_1 & L_2

$$\frac{|\lambda - 1|}{\sqrt{\frac{1}{a^2} + \frac{1}{b^2}}} = \frac{\sqrt{a^2 + b^2}}{2} \Rightarrow \lambda = 1 \pm \frac{a^2 + b^2}{2ab}$$

$$= \pm \left(\frac{a^2 + b^2}{2ab} \pm 1 \right)$$

$$= \pm \left(\frac{(a \pm b)^2}{2ab} \right)$$


$$PS^2 = PL^2$$

$$\left(x - \frac{a}{2} \right)^2 + \left(y - \frac{b}{2} \right)^2 = \left(\frac{x}{a} + \frac{y}{b} \pm \frac{(a \pm b)^2}{2ab} \right)^2$$

$$\frac{1}{a^2} + \frac{1}{b^2}$$

Ques

Find the locus of centres of a family of circles passing through the vertex of the parabola $y^2 = 4ax$ and cutting the parabola orthogonally at the other point of intersection.



$$(x-h)^2 + (y-k)^2 = (\sqrt{h^2+k^2})^2$$

$$x^2 + y^2 - 2hx - 2ky = 0.$$

$$(at^2, 2at) \rightarrow a^2t^4 + 4at^2 - 2h at^2 - 4ak t = 0.$$

$$\rightarrow at^3 + (4-2h)t - 4k = 0 \rightarrow t, \dots \quad \text{--- (2)}$$

$$\textcircled{2} - t \cdot \textcircled{1} \quad (4-2h)t - 4k + kt^2 - ht = 0.$$

$$kt^2 + (4-3h)t - 4k = 0 \quad \text{--- (3)}$$

$$tk = h + at^2$$

$$at^2 - kt + h = 0 \rightarrow t_1$$

$$\text{--- (1)} \rightarrow at^3 - kt^2 + ht$$

$$at^2 - kt + h = 0 \quad \text{--- ①}$$

$$kt^2 + (4 - 3h)t - 4k = 0 \quad \text{--- ②}$$

C.R.C.


$$(hk + 4ak)^2 = (a(4 - 3h) + k^2)(4k^2 - h(4 - 3h))$$

$$(h, k) \rightarrow (\lambda, \mu)$$

$$\mu^2(\lambda + 4a)^2 = (\mu^2 + 4a - 3a\lambda)(4\mu^2 - \lambda(4 - 3\lambda))$$

Que

If two chords drawn from the point $(4, 4)$ to the parabola $x^2 = 4y$ are divided by line $y = mx$ in the ratio $1 : 2$ then find the interval in which m lies.



$$C: (h, mh)$$

$$B: \left(\frac{3h-8}{3-2}, \frac{3mh-8}{3-2} \right)$$

$$\equiv (3h-8, 3mh-8)$$

$$x^2 = 4y$$

$$(3h-8)^2 = 4(3mh-8)$$

$$9h^2 + 64 - 48h = 12mh - 32$$

$$9h^2 - (48 + 12m)h + 96 = 0$$

$$D > 0$$

$$(m+4)^2 - 4 \cdot 9 \cdot 76 > 0$$

$$(m+4)^2 > 19$$

$$\Rightarrow |m+4| > \sqrt{19}$$

$$\Rightarrow m \in (-\infty, -4-\sqrt{19}) \cup (-4+\sqrt{19}, \infty)$$

Ans